

CLALLAM COUNTY ROAD DEPARTMENT

INTEGRATED WEED MANAGEMENT PLAN 2025



BIOLOGICAL



PHYSICAL



CHEMICAL



CULTURAL



PREVENTATIVE



POLLINATOR
FRIENDLY

The Cover: photos, from top Cinnabar Moth caterpillar (biological control agent) on tansy ragwort; staff manually removing tansy ragwort; selective herbicide treatment on ODT right of way; native planting crew of volunteers and CCNWB staff at ODT Berm on Old Olympic Highway; weed free stockpile to show prevention of weed seeds spreading; pollinator on goldenrod (*Solidago lepida*) in full bloom, one of the native species in our natural, pollinator-friendly, roadside revegetation projects.

Table of Contents

CLALLAM COUNTY INTEGRATED WEED MANAGEMENT PLAN	5
Introduction	5
Why Control Noxious and Invasive Weeds?	6
INTEGRATED WEED MANAGEMENT	7
Choosing Control Methods	7
The IWM Decision-Making Process	8
BIOLOGICAL WEED CONTROL	9
Roadside Application	9
PHYSICAL WEED CONTROL	11
Roadside Application	12
CHEMICAL WEED CONTROL.....	14
Roadside Application	14
OWNER WILL CONTROL AGREEMENT AND ADOPT-A-PATCH.....	18
CULTURAL WEED CONTROL	19
Roadside application	19
PREVENTATIVE WEED CONTROL.....	21
Roadside Application	21
2025 WORK PLAN.....	23
Identifying High Priority Weed Targets	24
2025 IWM Tasks.....	36
Application Locations.....	39
Selected Roads: Tables 4-6.....	40
East Clallam County Selected Roads	40
Central Clallam County Selected Roads.....	50
West Clallam County Selected Roads	56
County Pits	60
County Special Sites.....	62
Focus Area Maps of Selected Roads	66
Herbicide Product List.....	77
APPENDICES.....	78
Appendix A: Ordinance 923 - Integrated Weed Management	79
Appendix B: Non-Target Impacts and Risk Assessment, Herbicide Selection Process, Product List, and Application Methods	82
Appendix C: Sample Record Keeping Forms	107
Appendix D: Sample Press Release and Public Notice	111
Appendix E: Sample Herbicide Notice	112
Appendix F: Owner Will Control Packet.....	113
Appendix G: Adopt-a-Patch Permit.....	117
Appendix H: Roadside treatment activities 2024	118
Appendix I: References	133

List of Tables

	<u>Page</u>
Table 1. Insect biocontrol agents in Clallam County.....	10
Table 2. Known roadside weeds in Clallam County.	25
Table 3. Recommended control treatments for Clallam County roadside noxious weeds.....	31
Table 4. East Clallam County roads selected for herbicide treatment in 2024.....	41
Table 5. Central Clallam County roads selected for herbicide treatment in 2024.....	50
Table 6. West Clallam County roads selected for herbicide treatment in 2024.....	56
Table 7. County Pit Sites for treatment in 2024.....	60
Table 8. County Special Sites selected for treatment in 2024.....	62
Table 9. Species in Clallam County with potential for special management consideration.....	85
Table 10. Half-Maximum Application Rates Used in Risk Charts.....	87
Table 11. Toxicity reference values used to estimate risk	98
Table 12. Selected herbicide characteristics.....	104
Table 13. Adjuvants used to enhance herbicide effectiveness.....	105
Table 14. Historic roadside weed locations for Clallam County.....	118

List of Figures

	<u>Page</u>
Figure 1A. Himalayan blackberry monoculture.....	6
Figure 1B. Naturally sustainable roadside vegetation community.	6
Figure 2. Factors influencing IWM program.....	7
Figure 3. The IWM decision-making process.....	8
Figure 4. Risks to honey bees from direct spray or drift.....	89
Figure 5. Acute risks to aquatic invertebrates from first-flush runoff.....	90
Figure 6. Acute risks to fish from first-flush runoff.....	91
Figure 7. Acute risks to small mammals consuming contaminated fruit.....	92
Figure 8. Acute risks to small mammals consuming contaminated insects.....	93
Figure 9. Chronic risks to large mammals from consuming contaminated vegetation.....	94
Figure 10. Chronic risks to large birds from consuming contaminated vegetation.....	95
Figure 11. Acute risks to small birds consuming contaminated insects.....	96
Figure 12. Soil half-lives of herbicides in aquatic environments.....	101
Figure 13. Predicted concentrations from first flush runoff at half-maximum application rate...	102

CLALLAM COUNTY INTEGRATED WEED MANAGEMENT PLAN

INTRODUCTION

Clallam County is a major landholder subject to Washington State weed laws RCW 17.10 and WAC 16-750, which mandate the control of specific non-native, invasive “noxious” weeds. An integrated weed management plan is needed to help the County efficiently comply with its noxious weed control obligation on county roadsides and other lands under Road department management.

The primary responsibility of the Clallam County road system is safety in the transport of people, goods, and services. Roadside weed management is a unique element within a general road maintenance program. Effective weed management requires identifying the best mix of all control techniques as well as an understanding of how plant communities are part of a dynamic process. Healthy, self-sustaining plant communities better compete with weed pressure. The explicit goal of this plan is to shift roadside vegetation to natural, site appropriate plant communities. This includes using best management practices that support that shift, minimizing the need for all types of interventions, including the use of herbicide.

The IWM plan must be consistent with Clallam County’s long-term goals for its road system including environmental and public safety considerations. The IWM plan will strive for a balance of multiple, but compatible, goals, such as reducing maintenance costs for weed control over time while increasing environmental services. Other considerations will include protection and preservation of the natural environment, preserving and enhancing the scenic and habitat quality of the roadside and being a good neighbor to adjoining property owners.

This document serves as the strategic plan for managing non-native invasive plants that infest county rights-of-way. It contains information on priority weed locations and guidelines and procedures for best management practices in weed control. This plan is developed in compliance with Washington State Noxious Weed Law, Chapter 17.10 of the Revised Code of Washington, and modeled on the State of Washington’s Integrated Pest Management program as codified in Chapter 17.15 of the Revised Code of Washington. Specific County legislative direction upon which this plan is based is codified in Clallam County Code Ordinance 923 (Appendix A).

Roadside weed management is an evolving process and it is intended that this plan be annually evaluated and adapted over time based on input and technical updates from federal and state agencies, tribes, universities and local partners and stakeholders. It is essential that the results of control activities are monitored, evaluated and adjusted as necessary to maximize efficiency and effectiveness. Best Management Practices (BMPs) for each weed program element, in conjunction with continued research and education, will provide important information for ongoing Integrated Weed Management (IWM) treatments.

This plan encourages public involvement. “Owner Will Control” agreements, Adopt-A-Patch, and revegetation projects represent volunteer opportunities. Annual plans, treatment progress updates and year-end reports will be available to the public online at: <https://www.clallamcountywa.gov/821/Noxious-Weed-Control>

Please contact the County at the numbers listed below for questions or comments:

Christina St John, 360-417-2442

or

Joe Donisi, 360-417-2404

WHY CONTROL NOXIOUS AND INVASIVE WEEDS?

Noxious weeds impact native ecosystems by reducing biodiversity, altering hydrologic conditions, altering soil characteristics, changing fire intensity and frequency, modifying successional pathways, competing for pollinators, displacing rare plant species, serving as reservoirs of plant pathogens and by replacing complex native communities with simple non-native ones. Noxious weeds cause economic impacts. In general, noxious and invasive weeds are expensive to control and negatively impact agricultural and forestry production, property values, water flow and availability and recreation opportunities. It is estimated that invasive plants cause about \$123 billion in damages and losses to the U.S. economy annually (Harper-Lore, Johnson, and Skinner, 2007). Non-native weeds cause an estimated \$34 billion in losses to crops and pastures alone (Pimentel, McNair et al., 2001).

For these reasons, Washington State law requires the control of certain weed species. The purpose of the law is to limit economic loss and adverse effects to Washington's agricultural, natural, and human resources due to the presence and spread of noxious weeds in all terrestrial and aquatic areas in the state. The processes for regulation and control are defined in the Revised Code of Washington Chapter 17.10 and the Washington Administrative Code Chapter 16-750. All landowners, public and private, are required to control noxious weeds on lands they own.

Transportation rights-of-way are high priority locations for control of noxious weed species because they cross and link so many adjacent properties and land uses and can act as conduits for the spread of weeds. Weeds often appear first along road corridors.

Clallam County must be a responsible steward of County-owned land. It supports commerce and the economic viability of the agricultural community. The County also values environmental preservation. It has taken the lead on projects to restore ecosystem function. The County promotes tourism and recreational opportunities. All can be undermined by the spread of invasive plants. To meet its responsibilities, the County must ensure noxious and invasive weeds are effectively and efficiently controlled on its rights-of-way (Figure 1, A and B).



Figure 1A. The Himalayan Blackberry monoculture shown here, has overtaken the right of way after a road improvement project before native plants could be established and will outcompete all plants or trees until it is controlled.



Figure 1B. A low growing, naturally sustainable plant community is compatible with right-of-way goals while providing environmental services and quality habitat.

INTEGRATED WEED MANAGEMENT

Integrated Weed Management

Integrated Weed Management (IWM) is a coordinated decision-making process that uses the most appropriate weed management methods and strategies, along with a monitoring and evaluation system, to achieve roadside maintenance goals and objectives in an environmentally and economically sound manner. This includes assessing potential non-target impacts that may occur as a result and minimizing adverse effects through best management practices. The principles of IWM dictate that each weed problem is addressed from the perspective of all control options. The selected mix of control methods is the best treatment for the long-term stability of the plant community. Stable plant communities become established when the desired plants are not disturbed by the control program for the undesired plants. The physical design of the roadside environment coupled with the sporadic occurrence of noxious weeds imposes restrictions on the selection of control methods.




Figure 2. Factors influencing IWM program.

CHOOSING CONTROL METHODS

Weed control methods include biological, chemical, cultural, physical, and preventative. Each has its strengths and weaknesses influenced by regulations, environment, and economics (Figure 2). A consideration of potential non-target impacts also plays a role. (See Appendix B for risk assessments)

Biological (such as releasing insect agents) and physical methods (such as mowing) use fewer labor resources. These are best for managing and slowing the spread of, but not controlling or reducing, widely dispersed weed infestations. Mowing is non-selective and can spread weed seeds or other viable propagules. Biological agents can be extremely selective but require specific conditions. Both must be repeated indefinitely to suppress the weed population. Neither will eliminate populations of most weeds without using other techniques in combination.

Physical methods such as hand-pulling or digging are labor intensive but can effectively control or eliminate small weed infestations of limited distribution. While highly selective, such methods are unlikely to control deeply rooted weeds or weeds with spreading root systems.

Herbicides can effectively and selectively control all sizes and types of weed infestations with a small, but knowledgeable workforce. Careful attention must be paid to minimizing potential non-target effects and to follow all relevant regulations. See Appendix B for analysis of non-target impacts and risk assessment. Weather or site conditions can limit use.

Cultural and prevention practices are the most cost effective and efficient in the long term. These methods are more indirect and best used in tandem with the others. As current weed populations are eliminated, the goal is to shift control measures toward cultural practices such as use of native seed mixes and less disturbance of native shrub communities, as well as prevention practices such as weed-free material standards and cleaning equipment between job sites.

Use of the most effective method or combination of methods within an IWM decision-making framework will result in the greatest roadside service levels at the lowest life-cycle costs. Figure 3 demonstrates some of the feedback loops involved in an IWM strategy.

THE IWM DECISION-MAKING PROCESS




Figure 3. The continuing flow of monitoring, evaluation and adapting treatments occurring in an IWM program (adapted with permission from WSDOT area management plans).

BIOLOGICAL WEED CONTROL

Description

Methods which use living organisms to inhibit a host plant's ability to survive or reproduce are considered biological controls. Insects, diseases, and foraging animals, such as goats and cattle, are examples of biological control organisms. Biological methods are typically applied only when weed infestations are so well established that total eradication is not practical or possible.

General Use Considerations

Insect biocontrol agents are routinely inexpensive to maintain, but their populations lag behind the development of the weed population. Careful testing and screening are done before releasing insect agents to ensure they will not also attack native or other desirable plants. Insect predators are intended to be very weed-specific, though insects are not available for many weeds. They are usually part of regional programs of which the roadside right-of-way is an incidental beneficiary. Livestock grazing has the same effect as mowing; it removes the top growth without disturbing the roots so perennial plants re-grow as soon as grazing pressure is removed. Grazing animals can suppress desirable bee and butterfly forage, create bare ground or otherwise disturb the shoulder making it prone to re-infestation and erosion. Measures must be taken to ensure that animals do not destroy desirable vegetation on adjoining land. Contiguous acres are usually needed for successful biological control. Biological controls can reduce populations but can never result in eradication. The use of disease organisms as a form of biological control is still very limited.

ROADSIDE APPLICATION

There are several limitations and hazards associated with using grazing animals on Clallam roadsides. Most importantly, many of the noxious and invasive weeds targeted for control are sporadically dispersed along the road system and not easy to selectively target by grazing animals. Biological control is applicable where host weeds are present in dense or continuous colonies. All available insect agents have been released or are present for control of noxious weeds within Clallam County (Table 1).

Limitation

Grazing: *not considered for use at this time*

- Grazing animals on narrow right-of-way pose a hazard to motorists
- Insufficient grazing area
- Creates bare ground
- Targeted species are distributed in such a way that makes grazing inefficient and less selective

Insects:

- Insect agent unavailable for many weeds
- Non-contiguous infestation or insufficient host density
- Minimal disturbance is required for insect population to grow to an effective level; often conflicts with routine mowing schedule
- Will not eliminate weed populations, only suppress them

Table 1. Insect biocontrol agents in Clallam County

BIOLOGICAL AGENT	LATIN NAME	TARGET WEED	COMMENTS
Bindweed gall mite	<i>Aceria malherbae</i>	bindweed sp	Generally, for field bindweed, but experimental use for hedge bindweed; best for hot, dry sites; these are new releases
Bull thistle seed head gall fly	<i>Urophora stylata</i>	bull thistle	Seed feeder; not compatible with other control methods; may be present already
Canada thistle stem gall fly	<i>Urophora cardui</i>	Canada thistle	Metabolic sink, reducing vigor; not compatible with other control methods; three additional agents had been previously released (not by us!) for thistle control, but very detrimental to native thistle species.
Banded gall fly and knapweed seed head fly	<i>Urophora quadrifasciata</i> and <i>U. affinis</i>	knapweed, meadow	Flies often destroyed by seed weevils when both agents occur together
Lesser knapweed flower weevil and blunt nosed flower weevil	<i>Larinus minutus</i> and <i>L. obtusus</i>		Seed feeders
Knapweed root weevil	<i>Cyphocleonus achates</i> ,	spotted knapweed	Root weevil can be very effective, but limited distribution
Lesser knapweed flower weevil and blunt nosed flower weevil	<i>Larinus minutus</i> , and <i>L. obtusus</i>		Seed feeders
Sap-sucking psyllid	<i>Aphalara itadori</i>	knotweed sp.	Experimental releases only in WA
Defoliating hemlock moth	<i>Agonopterix alstroemeriana</i>	poison hemlock	Not effective, no longer distributed
Black-margined loosestrife beetle	<i>Galerucella californiensis</i>	purple loosestrife	Highly effective even in low density infestations, some non-target effects
Tansy flea beetle	<i>Longitarsus jacobaeae</i>	tansy ragwort	Best on rosettes and seedlings; poor survival in wet areas; a Swiss ecotype was released in the hopes of increased survival
Cinnabar moth	<i>Tyria jacobaeae</i>		A generalist that feeds on all plants in the <i>Senecio</i> genus, distribution not recommended - danger to native plants
Seed-feeding bruchid	<i>Exapion fuscirostre</i>	Scotch broom	Seed eaters
Seed-feeding weevil	<i>Bruchidius villosus</i>		
Klamath weed beetle	<i>Chrysolina quadrigemina</i>	St. John's wort	Feeds on ornamental and native species also; present; not well adapted to our climate

Application Guidelines

- Grazing may be possible under specific "Owner Will Control" option.
- Release additional viable insect agents should they become available.

Maintenance

None required at this time.

PHYSICAL WEED CONTROL

Description

Physical control includes both mechanized and manual methods. Mechanical methods use equipment to mow, cut, prune, scrape or cultivate in a manner which reduces, removes or prevents undesirable plant growth. A variety of machines are used in a roadside program, such as flail, reel, sickle, and rotary mowers, which come in different sizes, and graders, which are used to pull shoulders and remove sod buildup. Brush cutting is usually done with machines that are larger and heavier versions of rotary mowers. Mechanical methods are for larger scale general vegetation maintenance activities.

Manual methods include hand-held tools such as bladed weed-eaters, string trimmers, chain saws, brush hooks, hoes, and machetes; mechanical methods on a small scale, as well as grubbing and pulling weeds. Hand pulling is generally reserved for small or difficult to access sites or where greater selectivity is required. Repeat treatments are required for many species.

General Use Considerations

Mechanized equipment is typically used to non-selectively suppress undesirable or excessive vegetation growth on a large scale, not specifically to control weeds. Mechanical tools such as mowers do not affect the roots of plants and cut plants often resprout in greater numbers. This is particularly true of weedy biennial and perennial forbs or shrubs. Many weeds respond to mowing by shorter regrowth and producing seeds on stalks below the blade height. Properly timed or frequent mowing can delay or prevent seed development during a growing season, but improperly timed mowing results in spreading propagules over a broad area. Sod scalping causes erosion potential and creates sites for weed invasion.

Weed suppression by mowers is temporary and must be repeated to achieve the desired effect. Without specific guidelines, mowing is non-selective in its effect on the plant community. Many desirable native plants grow more slowly than their weedy, invasive cousins. Desired and undesired plants are continuously reduced to the same height, the same starting point, with each mowing. Some weeds are spread by the mowing operation. Stable plant communities, an expressed objective of the county's roadside program, are not retained under heavy mowing pressure. Unless carefully timed, close mowing may be disastrous for ground nesting birds, animals, and pollinator forage. Mowing also has a large carbon footprint in comparison to other control strategies and machinery can leak hydraulic fluid and shed other hazardous substances into ditches and other sensitive environments.

Special considerations for this management option are necessary due to exposure to hazards such as noise, sharp power equipment and road traffic. Extra alertness is necessary. Protection for eyes, ears, hands, legs, and feet is required when using these tools. Alternative mechanical methods such as steam or flame have been investigated. Both steam and flaming work by destroying top growth and are best used when plants are first germinating, not when well established. Both have little effect on roots. Additionally, steam and flaming pose significant hazards for the operator and the environment and are very costly. These methods are not being considered for inclusion currently.

Manual methods are commonly used for small infestations. This technique is effective in treating areas where obstructions prohibit mechanical methods. Hand pulling can be very selective and may be reserved for sites where extreme selectivity is critical and the infestation is small. Grubbing and hand pulling rely on moist soil and can be performed during inclement weather. These methods are labor intensive, slow, disturb the soil, and are usually expensive compared to other methods. As with other physical methods using mechanized equipment, manual treatments that do not extract all the roots of perennial plants will result in re-sprouting.

ROADSIDE APPLICATION

Physical control methods, both mechanical and manual, have been the preferred method for roadside vegetation management in Clallam County for many years. Mowing and brushing activities are an indispensable part of maintaining road safety by preventing line-of-sight obstructions, reducing fire hazard, preventing flooding, and ensuring bio-filtration of hazardous runoff. Reach mowing is the practice of clearing vegetation, primarily brush and small trees, from the right-of-way. Work is accomplished with a rotary or flail mowing head attached to an extendable boom mounted to a tractor. This practice includes ditches and intersections. Clearing undesirable brush and trees from ditches encourages the growth of desirable grasses. This helps maintain the bio-filtration function of grass, resulting in cleaner runoff water.

The road department strives to make one complete mowing pass per year: more at intersections or critical locations. Right-of-way mowing and brushing can occur from spring to early fall when shoulder vegetation is actively growing, and ongoing shoulder maintenance is required. Mowing does not normally reduce weed infestations but can provide temporary suppression. It is best used in close coordination with other weed control methods. In general, perennial weeds like Canada thistle must be mowed at least three times per season or the weeds are invigorated.

Under the IWM plan, road shop supervisors will work closely with the environmental coordinator, noxious weed control coordinator, and right-of-way weed crew lead and coordinate with the mowing crew to ensure that all work is performed in accordance with Endangered Species Act and water quality requirements and state weed laws. Critical areas have been identified ensuring that mowers will know which areas and locations require special consideration. Appropriate guidelines have been developed for these locations. Changes and updates are made as necessary.

The road department funds sheriff department-led Clallam Roadside Ecological Workers (C.R.E.W) provide a valuable manual workforce. Manual weed control activities will be incorporated into their assigned duties. C.R.E.W. or volunteer events will be the backbone of weed control activities that require a large labor force. They will be directed to work on large infestations of easily recognized weeds that can be effectively pulled such as flowering tansy ragwort and Scotch broom.

Limitation

- Mowing suppresses weed infestations; but does not control. Where control is desired, mowing is not recommended unless in combination with other control measures.
- Tough perennial weeds, especially those with extensive roots, are difficult to control using only physical means.
- Pulling or digging weeds is most effective when the ground is soft.

Application Guidelines

- Avoid close mowing of desirable, native vegetation. Limit back slope mowing as much as possible. Avoid mowing the back slope in critical areas.
- Resurvey mowed roads to locate weed regrowth.
- Do not mow knotweed infestations; mowing encourages re-sprouting, may spread fragments capable of producing viable plants, and makes other treatments less effective. Consult crews responsible for weed control recommendations.
- Manual methods may be applied where practical and conditions favorable.
- Digging should be limited to individual plants or very small infestations.
- Limit digging of perennial weeds or those with deeply spreading roots unless they are newly established.
- Pull and bag the heads of flowering plants. Dispose of appropriately.

Personnel

Mowing program operators are often rotated throughout the Road Maintenance crew. Each district has a different number of operators who mow, as time allows, between other road maintenance duties. We strive to aid the mower operators in their identification of noxious weeds; especially those that spread by propagules.

The road department funds C.R.E.W (as resources allow), which is comprised of up to five low-risk offenders overseen by a corrections officer. C.R.E.W performs various tasks as directed by the road department. Weed Board staff dig up minor regulated weed infestations as conditions and resources allow to help the County achieve compliance with law.

Training and Licensing

On-the-job training

Safety trainings, briefings

Plant identification and weed control training in conjunction with the Noxious Weed Control Program

Equipment

Various mechanical mowers and tractors with mowing attachments, weed whackers, chainsaws, weed wrenches, shovels, dandy diggers, and *hori-horis* (specialized digging tools).

Maintenance

- Regular maintenance and inspection of mowers and mower heads to minimize leaks or potential spills.
- Operators will be familiar with a spill prevention plan and carry spill kits.
- Ongoing training in critical areas issues for operators.
- Recurrent weed identification training for C.R.E.W.
- Ongoing improvements in equipment.

Performance Maintenance

- ◆ Number of weeds pulled
- ◆ Number of volunteer events
- ◆ Mowing – number of roadside pass miles. Pass miles count each shoulder mile mowed including those that are mowed more than once each year. The goal is to reduce this parameter while satisfying public, safety, and regulatory responsibilities.

CHEMICAL WEED CONTROL

Description

Herbicide applications only target specific noxious weeds or non-native and invasive species of special concern in our area that have been identified by the Clallam County Noxious Weed Board, state, or federal agencies. Applications are made with herbicides selected for their effectiveness on the weed being targeted and may be applied using backpack sprayers or other handheld equipment as determined appropriate by the site conditions and/or the target weed.

General Use Considerations

Herbicide applications are a less physically labor-intensive means of controlling large weed infestations. Herbicides are the most effective way to control deeply rooted, persistent weeds. Properly applied herbicides can suppress weed germination and allow desirable vegetation to flourish with minimal effort. However, herbicides may not be appropriate under certain site or weather conditions and require more complex decision making and staff training than most other control measures. In Washington all herbicides must undergo a registration process in addition to that required by the Environmental Protection Agency before they can be legally applied. Washington's pesticide laws may require an applicator to be licensed.

Choosing an herbicide application requires carefully considering the level of weed infestation, economic impacts, and human and ecologic consequences. When a chemical measure is chosen, optimal effect is achieved through proper herbicide selection, timely application, proper application method, and the use of the effective rate of herbicide.

Herbicide use may differ depending on the setting. Targeted roadside application, as included in this plan, is fundamentally different than that of many other types of applications where herbicides are the mainstay of weed control. Spot applications of herbicides in a noxious weed control program are often used to control individual plants, while in agricultural settings broadcast applications to entire fields are common. A limited number of chemicals are typically used for noxious weed management compared to those used in agriculture. In a successful weed management program, the amount of herbicide used on a particular site will decrease over time as the invasive plant population declines and sustainable, desirable plant replacement is supported. The potential for developing herbicide resistance is also significantly decreased by this approach.

An herbicide's potential risk is assessed by the Environmental Protection Agency before the product is registered for use. A clear understanding of the risk of using a particular herbicide requires knowledge of the toxicity of the herbicide as well as the likelihood of exposure. Toxicity is a measure of how harmful any chemical compound is. It can be measured in many ways and evaluated for many different biological systems. However, a chemical cannot have any effects on an organism without exposure. Because noxious weed management with herbicides necessarily introduces chemicals into the environment, the challenge is to estimate the amount of exposure (the dose) for humans and different types of animals, as well as non-target plants. The presence of an herbicide in the environment poses less risk if the exposure for non-target organisms is sufficiently low that it is unlikely to have a negative impact.

ROADSIDE APPLICATION

It is the explicit goal of this IWM plan to minimize the use of herbicides whenever practicable, while shifting roadside vegetation to natural, self-sustaining, site-appropriate plant communities. Activities that create bare ground while controlling weeds will be avoided, or be limited in duration, to prevent reinvasion by other weed species. Revegetation of bare ground with desirable plants will be promoted wherever opportunity exists.

Each species will have a Best Management Practice (BMP) specific to that species, developed and provided by the Clallam County Noxious Weed Program. Product label guidelines for timing and rates will be observed for best results. Herbicides may be used in conjunction with other practices, including biological and physical.

Most of the herbicides used in noxious weed control are of low toxicity; however, not all herbicides have equal impacts. For example, some may pose greater risks to aquatic life and are not approved for use in aquatic settings. Others have long-lasting pre-emergent herbicidal activity that may restrict plant emergence or growth for several months after treatment. In areas that are to be re-vegetated soon after treatment, these herbicides may not be the best choice if their residues remain biologically active in the soil after desirable plant species are seeded or transplanted.

Herbicide products chosen for this program are ones that maximize effectiveness, selectivity, and safety. Appendix B provides herbicide toxicity and possible exposure scenarios for wildlife. The analysis presented in the Cal-IPC document from which this information was reproduced, was based on the best available scientific data. Herbicide users are reminded of the need to keep in mind that risk analysis is a dynamic, ongoing process, as new data is generated on exposure potential and toxicity. Future studies or refined analyses may reveal risks that were previously unknown; alternatively, they may provide assurance that risks are lower than previously understood. Invasive weed managers must stay informed about the latest technical developments about the chemical and non-chemical strategies they use.

The way in which herbicides are applied can enhance efficiency and safety goals. Spot, foliar treatments with backpack sprayers or even more selective handheld equipment (such as wick applicators or injectors) will be the most used application method. Spot treatments can release or protect habitat for wildlife such as pollinators, songbirds, and small mammals. Spot treatments reduce potential for offsite chemical drift. No broadcast treatments with mechanized equipment are being considered.

The general treatment period for noxious weeds will be during the growing season when the weeds are in full leaf. Treating before bloom focuses on preventing seed production, treating after bloom focuses on herbicide translocation to the roots as the plant restores food levels in the roots. Late season treatments need to be timed so that green living leaf and stem growth is still present. Fall applications are effective for controlling germinating winter annuals, biennials in the rosette stage, and moving herbicide to the roots of established perennials.

All herbicides used by Clallam County are currently registered by the U.S. Environmental Protection Agency (EPA) and the Washington State Department of Agriculture (WSDA). Application of herbicides is in accordance with WSDA standards and chemical labels. County employees who apply the herbicides are licensed by WSDA or under the direct supervision of licensed personnel. In addition, these employees undergo continuous training to upgrade their expertise in the selection and safe application of herbicides. Herbicide labels, Safety Data Sheets (SDS), WSDA sensitive person list, a safety plan, and this document are kept in the office and in the weed control truck.

Record Keeping

Thorough record keeping is maintained on a WSDA approved form (Appendix C), per State requirement for all herbicide applications. The record includes information about the treatment including location, chemical used, weather conditions, and applicator comments. Citizen inquiries pertaining to herbicide applications are recorded and addressed.

Limitation

Herbicides should not be used:

- When heavy rainfall is imminent, winds exceed 10 mph or during other inclement weather conditions, such as heavy rainfall that is expected soon after an application.
- Where landowners have a current "Landowner Will Control" agreement
- Special management areas may have specific control practices or limitations.

Application Guidelines/Standard Operating Procedures

- Use only EPA and WSDA approved herbicides.
- All applications are conducted under direct supervision of licensed applicators.
- Observe strict compliance with product labels and with State and local regulations.
- Use personal protective equipment as directed on the herbicide product label.
- Carefully select products, rates, timing of application, and equipment to be used.
- Include marker dye to aid identification of treated areas.
- Follow all applicable notification protocols.
- Follow product label for use and storage.
- Apply only aquatically approved formulations within 25 feet of water.
- Treat only the noxious weed site.
- Minimize drift injury by not spraying when wind exceeds 10 mph.
- Use drift reduction agents or techniques as appropriate.
- Don't spray when drifting cannot be controlled.
- Avoid foliar application when measurable rainfall is imminent.
- Conduct mixing and loading operations in an area where a spill would not contaminate an aquatic site or well head.
- Do not rinse spray equipment near bodies of water or sources of potable water.
- Be aware and protective of people, working equipment, sensitive crops and gardens, apiaries, endangered species, water, and wells.
- Avoid direct applications to pollinators.
- Secure containers during transportation.
- Contain and clean up spills and request help as needed.
- Keep copy of product labels and SDSs in truck.
- Promptly respond to any public inquiries or direct them to the supervisor.
- Post treated areas and specify the duration of exclusion, if appropriate.
- Provide public educational information on the need for weed treatments.
- Coordinate weed management activities where joint use of a right-of-way exists.

Herbicide Notification Process

Our intent is to provide notice as far in advance as possible balanced with the ability to predict weather and scheduling. General notice is provided in early spring through a Press Release (Appendix D) provided to the local news media and Public Notice posted on the County website. Both include general vicinity of areas to be treated, reference to the IWM plan and how to obtain a copy, and information for entering into an Owner Will Control Agreement with Clallam County. Additionally, an Herbicide Notice (Appendix E) is posted at intersections within a quarter mile of each other or at the discretion of the applicator. The Herbicide Notice includes the herbicides to be used, target weed species, application date, and phone contact. Notices that are

pre-posted are re-dated with the actual date of application. Staff are trained and available to explain applications and answer onsite questions.

Staff

The IWM program will be staffed with a licensed supervisor and any additional employee will be licensed or operate under the direct supervision of the supervisor or licensed Noxious Weed Control Board staff.

Training and Licensing

Washington State Department of Agriculture Pesticide License “Public Operator”

Washington State University IPM Program Certification (Continuing Education)

Equipment

Equipment used: backpack sprayers, handheld spray bottles and loppers, wicker wipe applicators, EZ-Ject lance and injection guns for selected noxious weeds. A backpack sprayer is a self-contained unit (tank and pump) and is carried on the back of the applicator. The capacity of these sprayers is usually less than 5 gallons. The entire tank may be pressurized or only a small chamber that draws from the main tank. This equipment is useful for selective applications and spot treatments. Backpack sprayers are very adaptable to a wide range of nozzle configurations for treating foliage. The backpack sprayer is the major application device for roadside weed control in Clallam County.

Maintenance

- Regular maintenance and calibration of all spray equipment.
- Early detection of targeted weed infestations and ongoing site evaluations.
- Ongoing training of staff including yearly recertification credits.
- Ongoing improvement and updates of equipment and handling protocols.

Performance Measurements

- ◆ Number of projects completed and ratio of spot treatment to whole road treatment.
- ◆ Area of weeds controlled.
- ◆ Number of weed species controlled/encountered/eradicated.
- ◆ Public, interdepartmental, and agency weed control requests – number of requests, area of treatment, miles of road.
- ◆ Public satisfaction -- number of complaints (the lower the number, the better the performance)
- ◆ Survey goals: – area, number of miles inspected, number of roads in a four-year rotational cycle.
- ◆ Documentation, monitoring, evaluation, and reporting

OWNER WILL CONTROL AGREEMENT AND ADOPT-A-PATCH

Owner Will Control Agreement

Property owners have the option to keep the road right-of-way abutting their property weed free with or without herbicides. To do so, the property owner must enter into an Owner Will Control Agreement with the County and perform weed control as outlined in the Agreement.

When entering into an Owner Will Control agreement, property owners assume the County's responsibility under state laws RCW 17.10 and WAC 16-750 to control noxious weeds, which requires timely and often repeated control efforts during the growing season. The landowner would also assume any additional weed control responsibilities resulting from county policy.

Property owners participating in control agreements may also be interested in assisting with cultural control enhancements consistent with long-term roadside weed control goals. Such opportunities will be pursued as program resources and voluntary participation allow.

A sample Owner Will Control packet is included in Appendix F.

Adopt-A-Patch

The public may "Adopt-A-Patch" consisting of weeds infestations that may be controlled manually. These types of locations will be annually identified, compiled, and posted online before seasonal county treatments commence. Those who wish to adopt a patch must enter into a Control Agreement (obtain a special use permit) with the County, which will outline control responsibilities and the deadline for completion.

An adoption deadline will be posted along with Adopt-A-Patch locations to avoid scheduling conflicts with County control personnel. Permits granting permission to control weeds on County-owned or managed lands, safety training and tips, as well as any other conditions required by county policy will be included in the Control Agreement packet.

A sample Adopt-a-Patch packet is included in Appendix G.

CULTURAL WEED CONTROL

Description

Techniques that benefit the development and health of desirable, competitive plant communities are considered cultural weed control methods. Cultural methods, along with prevention, are the preferred method of weed control wherever possible. Examples include the use of mulch and soil amendments that improve soil fertility to stimulate growth of desired species or to alter soil pH to discourage undesired plants. Cultural weed control includes the planting or seeding of desirable species. Effective use of cultural methods must be conducted in close coordination with efforts to eliminate existing noxious weed sites.

General Use Considerations

Cultural control methods encourage natural, self-sustaining, site-appropriate plant communities to develop in the long term. Native plant materials are preferred because once established on appropriate sites they require few additional inputs to thrive and self-perpetuate. In addition to low maintenance, well established native plantings provide many environmental services, such as erosion control, bio-filtration, pollinator, and animal habitat. Native plantings have the potential to prevent undesirable weeds from becoming established by two mechanisms, competition, and allelopathy. Competition is the interaction between plants for site resources such as space, nutrients, moisture, and light. Allelopathy occurs when one plant produces chemicals which inhibit the establishment and growth of others. The composition of plant communities on the roadside is likely to be a result of both mechanisms. Native wildflowers provide forage for pollinators and are aesthetically pleasing, while occupying the site to prevent or retard invasion by undesired noxious weeds.

ROADSIDE APPLICATION

The long-term goal of this plan is to programmatically incorporate cultural practices into overall roadside management practices. Possible opportunities include new construction, shoulders and ditches, locations under "Owner Will Control" agreements, post-treatment, and other county land such as pits, trails, and parks.

Activities that enhance or create native or self-sustaining plant communities should be applied as broadly as possible. Cultural practices are best applied to disturbed or bare ground or after weed treatments have occurred. Controlling the noxious weeds may release native roadside plants but more active measures may be required. Clallam County is committed to developing and obtaining locally sourced native seed mixes and plant material as available.

Activities to improve site conditions such as weed control, mulching or soil amendments to increase successful desirable plant establishment will be considered as resources and materials are available. Such activities cannot interfere or conflict with the primary use and safety of county rights-of-way.

Limitation

- Revegetating activities must be postponed until weed infestation is adequately controlled.
- Plant selection must not conflict with roadside safety and maintenance considerations, public or animal health, and adjacent land use or values.
- Roadsides are a harsh environment for establishing many desirable plant species; amending soil and providing supplemental water may not be a viable option in many cases.

Application Guidelines

- Use native species wherever possible.
- Blend with adjacent landscaping.
- Choose low growing plants for foreslope that require less mowing.

- Utilize weed-free, chipped materials on site to suppress weeds. Weedy brush may only be chipped and left on site if it is in early growth stages and has no ripe seed.

Staff

WSU Master Gardener Coordinator
Road Department Volunteer Coordinator
Volunteers
Weed Crew

Training and Licensing

No licensing required
Native plant identification, biology, and habitat needs
Planting techniques
Use of GPS equipment

Equipment

Handheld GPS, hand tools

Maintenance

Maintain/evaluate sites for first five years
Monitor periodically thereafter

Performance Measurements

- ◆ Cooperative relationships with outside entities developed and maintained
- ◆ Planning documents developed
- ◆ Volunteer participation
- ◆ Area vegetated with native, self-sustaining plant community
- ◆ Maintenance costs are reduced over time

PREVENTATIVE WEED CONTROL

Description

Preventative weed control refers to any control method that aims to reduce or prevent weeds from being established. Examples of preventative weed control would be using certified weed free materials such as road and shoulder base rock, gravel, straw, soil, or mulch material for construction and maintenance activities, and making sure equipment is cleaned before moving from one location to another.

General Use Consideration

Prevention is, by far, the most environmentally and cost-effective control strategy. In addition to the above-mentioned best management practices, prevention includes actively surveying for and eradicating new invaders or small, newly discovered infestations as they are encountered.

ROADSIDE APPLICATION

Prevention is a top priority for this plan. Using certified weed-free materials whenever possible reduces or prevents introducing new weeds and avoiding soil disturbance helps prevent creating an environment vulnerable to invasion. Road designs that are easier to maintain and incorporate weed prevention features can be very cost effective in the long term. Weed Board staff is available to advise and provide technical assistance to Road Department engineers at all stages in road construction.

Weed Free Materials. Since prevention is the foundation of noxious weed control, prevention should start with certified weed-free seeds, mulches, soil, and gravel. The North American Invasive Species Management Association has certification standards that involve inspection of sources and sites to determine they do not contain seeds or plant parts of invasive weeds. Inspection includes, but is not limited to, surrounding ditches, topsoil piles, gravel/sand piles and pits, fence rows, roads, easements, rights-of-way, working areas, storage areas, and a buffer zone surrounding the area. Washington subscribes to these certification standards and Clallam County will apply these standards as widely as possible.

Clean Equipment. An important part of prevention is to not carry noxious weed seeds or plant parts from site to site. Before moving from or to a construction or maintenance project, clean the equipment. Remove hanging debris; wash off mud. Ensure that associated vehicles and crew are similarly inspected. These actions help stop the movement of weeds along the roadside corridor. Clean hand tools, boots, and clothing as well.

Avoid Bare Ground. One of the problems of mechanical tools is scalping the soil. Whether it's by a grader, a mower, a bush hog, or a string trimmer, bare ground creates openings for the establishment of invasive weeds. Historically, when roadsides were disturbed, native plants from the soil seed bank or undisturbed adjacent land provided the seed source for the new native plants. In today's world, with human disturbances and inadvertent plant introductions, aggressive invasive plants are ready to occupy the available site. Covering bare ground with weed-free materials or seeding with desirable seed mix as soon as possible will suppress weed germination.

Design Controls. Adding new engineering standards that require less maintenance, such as favorable slope gradient, extending chip-seal edge, and incorporating native plantings in construction planning all help to prevent weed invasion.

EDRR. Eradication is a very realistic objective in the early stage of noxious weed establishment. Detecting new invaders or small weed patches and eliminating them at an early stage prevents costly

intervention later. This form of prevention, called Early Detection, Rapid Response (EDRR), is a preferred strategy for this program.

Both county employees and the public can be an important part of the EDRR process. The components of EDRR are: 1) detection reporting, 2) identification confirmation, 3) rapid assessment, 4) program planning, and 5) rapid response. The public will be encouraged to report suspicious plants, or new weed locations. The Clallam County noxious weed board has the resources to confirm the identity of suspicious weeds.

Limitation

- Roadways are exposed to all manner of weed pressure and completely preventing transportation and introduction of invasive plants is not possible.
- Routine maintenance activities will create some amount of bare ground.
- This control measure does not deal with established weed infestations.

Application Guidelines

- Incorporate prevention strategies programmatically into all aspects of planning and executing weed control activities and road maintenance.
- Develop native plant materials so that native seeds of desired plants are readily available.
- Limit activities that create bare ground.
- Where disturbance is expected, plan to revegetate with site appropriate plants. Identify the most favorable conditions for establishment.
- Inspect, evaluate weed invasion risk, and treat appropriately in response to emergency disturbances such as fire and flooding.
- Adopt a monitoring schedule to detect the presence of new invaders along roadsides or weed invasion of new construction.
- Incorporate EDRR strategy

Personnel

WSU Master Gardener Coordinator
Volunteers
Noxious Weed Control Board Staff

Training and Licensing

Cooperative training with WSDOT
Annual prevention and weed identification programs conducted by the Weed Board

Equipment

None determined at this time

Performance Measurement

- ◆ A higher percentage of weed sites are small.
- ◆ Program costs are reduced over time.

2025 WORK PLAN

2025 Work Plan

The focus of this work plan is the control of state-listed noxious weeds and invasive, non-native weeds of special concern to Clallam County rights-of-way. The integral precept of this IWM plan is that all control techniques are potentially applicable to the solution of the problem. A project list will be posted online and updated as treatments occur to keep the public informed.

With more than five hundred miles of county roads there are a variety of weed problems as well as control opportunities. Biological controls will continue to operate on roadsides through releases made elsewhere in the county. Additional releases will be made if new insect controls become available and are compatible with routine maintenance activities such as mowing, which is non-specific to and independent of weed control.

Physical controls will continue to be applied across the road system where effective and as resources allow. Volunteers will be recruited for various projects, especially where adjacent infestations threaten county assets.

Chemical control is an important tool that is needed for specific weed problems. Great care has been taken in choosing which herbicides may be applied, and additional safeguards are included by ensuring only targeted, hand applications are allowed. Herbicide use is limited to specific county roads and noxious weed locations which are listed in this plan based on past surveys and on-going treatment locations.

Most importantly, cultural and preventative controls will be applied programmatically to sustain the progress made by all the above-mentioned control methods. Combined, these management practices will move us towards achieving a low maintenance, naturally stable plant community.

High priority weed targets are identified and control options for an array of roadside weeds are summarized in the tables below. These are followed by specific tasks necessary to implement the 2025 work plan. Tasks are itemized under separate category headings. While listed separately, the tasks represent the best mix of control options chosen to address specific weed problems. The complete set of tasks is carefully designed to be implemented in tandem, not independently.

IDENTIFYING HIGH PRIORITY WEED TARGETS

Table 2 contains many of the known roadside noxious weeds for Clallam County. The table is arranged to show which weeds are the highest priority for control based on potential economic or environmental impacts and feasibility for control. The list is not comprehensive and will change as conditions change.

"Plant status" indicates one of several categories: a **noxious weed** (a prioritized legal designation including Class A, Class B and Class C weeds where control may be required under state law), a **non-native, invasive** plant capable of causing economic or environmental impacts, but not listed by the state, and **weedy**, so prevalent that it is generally considered naturalized or an aesthetic nuisance. Infestations of invasive, non-native species are more easily eliminated before they become established.

To be most efficient when deciding treatment priorities, where known, weeds are characterized as widespread or rare. The following abbreviations are used in the "status" column in Table 2:

ISSC = Invasive **S**pecies of **S**pecial **C**oncern

NCR = Noxious, **C**ontrol **R**equired

NR = Noxious, **R**are

NW = Noxious, **W**idespread

WR = **W**eedy, **R**are

WW = **W**eedy, **W**idespread

Weeds are assigned to a "category" based on information in the "status" column. Weeds are categorized as follows:

Category 1 weeds are Class A, B designate, and selected B or C noxious weeds, additional noxious weeds and invasive species of special concern that are very limited in distribution, and newly discovered invaders that were previously unknown in the county (EDRR - early detection, rapid response). Category 1 weeds are the *highest priority* for control.

Category 2 weeds are noxious weeds that are widespread, but of particular concern to the public or an affected public entity. Category 2 weed infestations will be added to the annual work plan to methodically reduce widespread weeds over time and to accommodate requests.

Category 3 weeds are those that are so widespread they are generally considered naturalized or a nuisance. These weeds are tolerated. Control is not considered feasible.

Table 2. Known roadside weeds in Clallam County

COMMON NAME	SCIENTIFIC NAME	4- LETTER WEED CODE	CATEGORY	STATUS	THREAT
Category 1 Weeds					
alyssum, hoary	<i>Berteroa incana</i>	BEIN	1	NCR	Aggressive invader in fields of forage crops; toxic to horses
blueweed	<i>Echium vulgare</i>	ECVU	1	NR	Invades pastures and meadows and outcompetes desirable plants; only known from one NPS road
brome, false	<i>Brachypodium sylvaticum</i>	BRSL	1	NCR	Aggressive bunch grass that is highly invasive in shaded woodlands, open prairies, and grasslands; only known from one location on FS road
chervil, wild	<i>Anthriscus sylvestris</i>	ANSY	1	NCR	Will grow in many soil types and light levels; quickly forms monocultures and poses a threat to agriculture and natural areas
cinquefoil, sulfur	<i>Potentilla recta</i>	PORE	1	NCR	Not readily grazed by livestock and wildlife; dense stands exclude other species
comfrey, common	<i>Symphytum officinale</i>	SYOF	1	ISSC	Used medicinally for poultices; liver damage when ingested; can form dense stands; difficult to control once established
fennel, common	<i>Foeniculum vulgare</i>	FOVU	1	NCR	Dense stands exclude native vegetation; outcompetes desirable forbs and grasses in pastures and grasslands
hairy whitetop	<i>Lepidium appelianum</i>	LEAP	1	NR	Monocultures displace desirable plants; unpalatable; can be toxic to cattle
hawkweed, European	<i>Hieracium sabaudum</i>	HISA	1	NCR	Dense stands exclude other species; bitter and unpalatable, little forage for livestock and wildlife
hawkweed, orange	<i>Hieracium aurantiacum</i>	HIAU	1	NCR	Dense stands exclude other species; bitter and unpalatable, little forage for livestock and wildlife
hawkweed, yellow	<i>Hieracium caespitosum</i>	HICA	1	NCR	Dense stands exclude other species; bitter and unpalatable, little forage for livestock and wildlife
hogweed, giant*	<i>Heracleum mantegazzianum</i>	HEMA	1	NCR	Skin contact with sap causes severe dermatitis and blistering on people and animals
Italian arum	<i>Arum italica</i>	ARIT	1	NR	Toxic; forms dense monocultures; difficult to control once established; rare in Clallam County

COMMON NAME	SCIENTIFIC NAME	4- LETTER WEED CODE	CATEGORY	STATUS	THREAT
Jewelweed, spotted	<i>Impatiens capensis</i>	IMCA	1	NCR	Invades riparian areas and wetlands, roadsides; only found on private and FS roads
knapweed, diffuse	<i>Centaurea diffusa</i>	CEDI	1	NCR	Spreads seed by tumbling; prickly flower heads; unpalatable after early spring
knapweed, meadow	<i>Centaurea x gerstlaueri</i>	CEMO	1	NCR	Outcompetes pasture species; degrades wildlife habitat; interferes with agriculture
knapweed, mountain	<i>Centaurea montana</i>	CEMO2	1	ISSC	Dense stands exclude native vegetation; out-competes natural forage species
knapweed, spotted	<i>Centaurea stoebe</i>	CEST	1	NCR	Allelopathic plant that can inhibit the germination of grasses; forms dense stands that exclude desired plants and wildlife
knotweed, Bohemian	<i>Fallopia X Bohemian</i> (<i>Polygonum bohemicum</i>)	POBO	1	NCR	Easily spreads by disturbance; dense colonies eliminate other plant species and can degrade fish habitat; causes structural damage
knotweed, giant	<i>Fallopia sachalinensis</i> (<i>Polygonum sachalinense</i>)	POSA	1	NCR	Easily spreads by disturbance; dense colonies eliminate other plant species degrades fish habitat; structural damage to human structures
knotweed, Japanese	<i>Fallopia japonica</i> (<i>Polygonum cuspidatum</i>)	POJA	1	NCR	Easily spreads by disturbance; dense colonies eliminate other plant species and can degrade fish habitat; causes structural damage to human structures
kochia	<i>Bassia scoparia</i>	BASC	1	NCR	Serious agriculture and pasture pest; Annual that quickly forms monocultures in disturbed, open sites; flammable; toxic to livestock; only found on Hwy 101 currently
loosestrife, purple*	<i>Lythrum salicaria</i>	LYSA	1	NCR	Dense stands eliminate other plant species; poor palatability; degrades wildlife habitat and hunting and fishing areas.
lupine, tree	<i>Lupinus arboreus</i>	LUAR	1	WR	Can hybridize with native lupine; forms dense monocultures; drought tolerant, likes disturbed areas
moleplant	<i>Euphorbia lathyris</i>	EULA	1	ISSC	Latex causes severe skin irritation and can be fatal if ingested; spreads rapidly
nightshade, bittersweet	<i>Solanum dulcamara</i>	SODU	1	NR	Aggressive and vine like, grows in disturbed lands and pastures; toxic to livestock

COMMON NAME	SCIENTIFIC NAME	4- LETTER WEED CODE	CATEGORY	STATUS	THREAT
old man's beard	<i>Clematis vitalba</i>	CLVI	1	NR	Climbing growth smothers other plants, even trees
poison hemlock	<i>Conium maculatum</i>	COMA	1	NCR	Highly toxic to humans and animals; all parts of the plant are toxic; severe birth defects
ribbon grass	<i>Phalaris arundinacea</i> cultivar	PHAR2	1	NR	Aggressive invader displaces other plants in wet sites; an ornamental form of reed canary grass; may also be used as a source for psychedelic drugs
rush skeletonweed	<i>Chondrilla juncea</i>	CHJU	1	NR	Aggressive invader; displaces native plant communities; only known from one historical site on Hwy 101
sowthistle, perennial	<i>Sonchus arvensis</i>	SOAR	1	NR	Aggressive invader; outcompetes desirable forbs and grasses in pastures; limited distribution in Clallam County
spurge laurel	<i>Daphne laureola</i>	DALA	1	NR	Toxic to humans and animals; contact with plants can cause dermatitis
tansy, common	<i>Tanacetum vulgare</i>	TAVU	1	NCR	Dense stands degrade forage value; toxicity issues for humans and livestock
tansy ragwort	<i>Jacobaea vulgaris</i> (<i>Senecio jacobaea</i>)	JAVU	1	NCR	Poisonous to horses, cattle, and pigs; animals grazing tansy can produce tainted milk, may result in toxic residue in honey
teasel, common	<i>Dipsacus fullonum</i>	DIFU	1	NCR	Forms dense stands of prickly, unpalatable plants; degrades habitat and reduces accessibility
thistle, Italian	<i>Carduus pycnocephalus</i>	CAPY	1	NCR	Forms dense stands of prickly, unpalatable plants; potential roadside fire hazard
toadflax, Dalmatian	<i>Linaria dalmatica</i> ssp. <i>dalmatica</i>	LIDA	1	NCR	Invades pastures and meadows, outcompeting desirable plants; only 1 known infestation on Hwy 101
wild basil savory	<i>Clinopidium vulgare</i>	CLVU	1	NCR	Aggressive invader, outcompetes native understory
willow herb, hairy	<i>Epilobium hirsutum</i>	EPHI	1	NCR	Semi aquatic in wetlands, ditches, streambank; dense growth can outcompete native vegetation
wormwood, absinth	<i>Artemisia absinthium</i>	ARAB	1	NR	Aggressive invader, will outcompete desirable forbs and grasses in pastures, fields and native grasslands

COMMON NAME	SCIENTIFIC NAME	4- LETTER WEED CODE	CATEGORY	STATUS	THREAT
yellow archangel	<i>Lamiastrum galeobdolon</i>	LAGA	1	NCR	Aggressive invader, competes understory species, degrades wildlife habitat
Category 2 Weeds					
bamboo, spp	<i>Phyllostachys aurea</i>	PHAU	2	NR	Bamboo species form dense thickets and roots; jeopardize line of site and asphalt on county roads; causes structural damage to human structures
barberry, common	<i>Berberis vulgaris</i>	BEVU	2	NR	Can form undesirable thorny thickets that can crowd out native species.
bindweed, field	<i>Convolvulus arvensis</i>	COAR	2	NR	Outcompetes native vegetation and reduces crop yields; climbs and forms dense tangled mats
blackberry, cutleaf	<i>Rubus laciniatus</i>	RULA	2	NW	Dense canopies crowd out native species; impenetrable barrier
blackberry, Himalayan	<i>Rubus armeniacus</i>	RUAR	2	NW	Dense canopies crowd out native species; impenetrable barrier
broom, Scotch	<i>Cytisus scoparius</i>	CYSC	2	NW	Forms dense stands; unpalatable; interferes with forest regeneration; fire hazard; scent can exacerbate human grass allergies; seeds are toxic to horses and livestock
burdock, common	<i>Arctium minus</i>	ARMI	2	WR	Forms large rosettes; hooked spines on seeds become entangled in fur of animals
butterfly bush	<i>Buddleia davidii</i>	BUDA	2	NR	Invades natural areas; crowds out native vegetation in riparian areas and interferes with natural succession
canarygrass, reed	<i>Phalaris arundinacea</i>	PHAR	2	NW	Unpalatable unless young, forms dense stands that crowd out native plants; especially difficult to control; serious wetland invader; stops process of succession in riparian sites, impedes tree seedling
cheatgrass or downy brome	<i>Bromus tectorum</i>	BRTE	2	ISSC	Depletes soil moisture in early spring; fire hazard in summer; known to occur in Clallam County, but not on roadsides; will be treated under EDRR protocol if observed
chicory, common	<i>Cichorium intybus</i>	CIIN	2	ISSC	Aggressive invader along roadsides, abandoned lots, lawns and overgrazed pastures

COMMON NAME	SCIENTIFIC NAME	4- LETTER WEED CODE	CATEGORY	STATUS	THREAT
common mullein	<i>Verbascum thapsus</i>	VETH	2	WR	Formerly on State Weed list; unpalatable, not widespread
hawthorn, English	<i>Crataegus monogyna</i>	CRMO	2	NR	Carried by birds into forests and open fields where it can form dense, thorny thickets that outcompete native species
herb Robert	<i>Geranium robertianum</i>	GERO	2	NW	Rapid spreading; displaces native herbaceous plants; allelopathic, inhibits the germination of small-seeded forbs in forest understory
iris, yellow flag	<i>Iris pseudacorus</i>	IRPS	2	NR	Toxic to humans and animals; displaces vegetation at wet margins of ditches, ponds, and lakes; plant resins can cause skin irritation in humans
Ivy, English	<i>Hedera helix</i> varieties	HEHE	2	NW	Forms dense carpets in understory; climbs and can kill trees or make them more prone to topple during storms; sap can cause allergic reactions and plant is toxic to humans and cattle
mountain ash, European	<i>Sorbus aucuparia</i>	SOAU	2	ISSC	Escaped ornamental that invades natural forests and grows rapidly; can quickly begin to outcompete native vegetation; seeds spread readily by birds
Mustard, wild	<i>Brassica spp.</i> (or <i>Sinapis arvensis</i>)	BRSP	3	WW	Common weed in fields, roadsides and abandoned areas; can interbreed with domestic Brassicas and contaminate seed crops
nightshade, hairy	<i>Solanum physalifolium</i>	SOPHY	2	NR	Thrives in degraded pastures and disturbed areas, poisonous to humans and livestock
peavine, everlasting	<i>Lathyrus latifolius</i>	LALA	2	ISSC	Forms dense thickets; seeds can be toxic to livestock; seriously interferes with forest regeneration where it invades from edges of timber units
periwinkle, large	<i>Vinca major</i>	VIMA	2	WW	Dominates in forest understories and displaces native vegetation
St John's wort, common	<i>Hypericum perforatum</i>	HYPE	3	NW	Causes photosensitization when grazed; toxic at all stages of growth
thistle, bull	<i>Cirsium vulgare</i>	CIVU	2	NW	Aggressive competitor; unpalatable for cattle; decreases forage

COMMON NAME	SCIENTIFIC NAME	4- LETTER WEED CODE	CATEGORY	STATUS	THREAT
thistle, Canada	<i>Cirsium arvense</i>	CIAR	2	NW	Aggressive competitor, unpalatable; decreases forage; host species for several agricultural pests
toadflax, yellow	<i>Linaria vulgaris</i>	LIVU	2	NR	Invades rangelands and cultivated fields; reduces yields; only known on Hwy 101

Category 3 Weeds					
buttercup, creeping	<i>Ranunculus repens</i>	RARE	3	WW	Depletes wet soil potassium; grows 40 sq ft/year; toxic to animals
carrot, wild	<i>Daucus carota</i>	DACA	3	NW	Damages agricultural commodity as it may cross pollinates with domestic carrot, seriously degrading the quality of commercial carrot seed production
catsear, common	<i>Hypochaeris radicata</i>	HYRA	3	NW	Crowds out palatable forage species; thrives in highly disturbed areas
chervil, burr	<i>Anthriscus caucalis</i>	ANCA	3	WW	Aggressive and can form monocultures; displaces native vegetation in pastures and along ROW's.
daisy, oxeye	<i>Leucanthemum vulgare</i>	LEVU	3	NW	Livestock avoid grazing; milk from dairy cows has unpleasant flavor
dandelion, common	<i>Taraxacum officinale</i>	TAOF	3	WW	Dense circular mats that crowd out desirable species
English holly	<i>Ilex aquifolium</i>	ILAQ	3	WW	Dense thickets can dominate shrub layer and suppress desirable vegetation
foxglove	<i>Digitalis purpurea</i>	DIPU	3	WW	Can be toxic to livestock; spreads aggressively in disturbed areas
orchard grass and other pasture grasses	<i>Dactylis glomerata</i>	DAGL	3	WW	Outcompetes and suppresses native vegetation including pasture grasses

*No active sites, but previously documented

Table 3 shows general guidelines for year-round treatments of the listed noxious weeds and invasive weed species of special concern. It is intended as a basic reference framework from which decisions are made for weed treatments from available options. Seasonal variables are considered and addressed as they become evident. The Clallam County Noxious Weed List or species that appear on county right-of-way may adjust as necessary.

Table 3: Recommended control treatments for Clallam County roadside noxious weeds

NOXIOUS WEED	FALL	WINTER	SPRING	SUMMER
Category 1 Weeds				
alyssum, hoary	Foliar herbicide treatment	Manual removal/digging; Foliar herbicide treatment	Manual removal/digging; foliar herbicide treatment before bolting	Manual removal/digging
blueweed	Manual removal/digging; Foliar herbicide treatment	No action	Foliar herbicide treatment	Manual removal/digging; Foliar herbicide treatment
brome, false	Manual removal; foliar herbicide treatment	Manual removal/digging	Foliar herbicide treatment	Manual removal/digging
chervil, wild	Foliar herbicide treatment for fall regrowth	Foliar herbicide application in late winter	Foliar herbicide treatment (rosette stage); manual removal of flowering plants	Manual removal of flowering plants
cinquefoil, sulfur	Removal of seed heads	Plants die back - no action	Foliar herbicide treatment of rosettes	Removal of flowering stalks
comfrey, common	Foliar herbicide treatment	Plants die back - no action	Foliar herbicide treatment	Foliar herbicide treatment
fennel, common	Removal seed heads	No action	foliar herbicide treatment before flowering	Cut flowering stalks
hairy whitetop	Foliar herbicide treatments	Plants die back - no action	Foliar herbicide treatments	Foliar herbicide treatments
Hawkweed, European	No action	Plants die back- no action	Foliar herbicide treatments	Foliar herbicide treatments; removal of flowering plants
hawkweed, orange or yellow	No action	No action	Foliar herbicide treatment	Foliar herbicide treatment; cut and remove flowers

NOXIOUS WEED	FALL	WINTER	SPRING	SUMMER
hogweed, giant	Manual removal/ digging – taproot should be cut 3-4” below soil level	Plants die back - no action	Foliar herbicide treatment; Manual removal/ digging – taproot should be cut 3-4” below soil level	Manual removal/ digging – taproot should be cut 3-4” below soil level
Italian arum	Foliar herbicide treatment; manual removal of small infestations	Foliar herbicide treatment; manual removal of small infestations	Foliar herbicide treatment; manual removal of small infestations	Plants die back – no action
jewelweed, spotted	Manual removal	No action – summer annual	Foliar herbicide treatment	Manual removal
knapweed species	Foliar herbicide treatment	Manual removal/ digging - small plants only	Foliar herbicide treatment	Foliar herbicide treatment; cut flowering stalks
knotweed species	Foliar herbicide treatment or injection	Plants die back - no action	No action	Foliar spray or injection (late summer)
kochia	Manual removal	No action – summer annual	Manual removal; foliar herbicide treatment	Manual removal
loosestrife, purple	Foliar herbicide treatment	Plants die back - no action	Foliar herbicide treatment	Foliar herbicide treatment
lupine, tree	Foliar herbicide treatment; manual removal	Manual removal	Foliar herbicide treatment	Foliar herbicide treatment; remove flowering stalks
moleplant	Foliar herbicide treatments if actively growing	Manual removal/ digging	Foliar herbicide treatments	Foliar herbicide treatments
nightshade, bittersweet	No action	No action	Foliar herbicide treatments	No action
old man’s beard	Foliar herbicide treatment if prostrate; basal stem treatment	Basal stem treatment	Foliar herbicide treatment if prostrate; basal stem treatment	Foliar herbicide treatment if prostrate; basal stem treatment
poison hemlock	Manual removal of flowering plants; herbicide treatment for fall regrowth for rosettes	Foliar herbicide treatment – rosette stage (late winter)	Foliar herbicide treatment of rosettes; Manual removal of flowering plants	Foliar herbicide treatment of rosettes; Manual removal of flowering plants;
ribbon grass	Foliar herbicide treatment	Plants die back - no actions	Foliar herbicide treatment	Cut flowering heads

NOXIOUS WEED	FALL	WINTER	SPRING	SUMMER
rush skeletonweed	Foliar herbicide treatment	No action	Foliar herbicide treatment	Cut flowering heads; foliar herbicide treatment
sowthistle, perennial	Cut flowering heads; foliar herbicide treatment	Plants die back - no actions	Foliar herbicide treatment	Cut flowering heads
spurge laurel	Foliar herbicide treatment; cut stump	Manual removal/ digging; cut stump or basal bark	Foliar herbicide treatments; cut stump	Foliar herbicide treatments; cut stump
tansy, common	Foliar herbicide treatment; cut flowering heads	Plants die back - no action	Foliar herbicide treatment	Cut flowering heads
tansy ragwort	Manual removal of flowering plants; foliar herbicide treatment of rosettes for fall regrowth	No action	Foliar herbicide treatment (rosette stage)	Manual removal of flowering plants; foliar herbicide treatment of rosettes
teasel, common	Foliar herbicide treatment of rosettes	No action	Foliar herbicide treatment	Manual removal of flowering plants; foliar herbicide treatment of rosettes
thistle, Italian	No action	Manual removal/ digging; foliar herbicide treatment	Manual removal; foliar herbicide treatment	No action – winter annual
toadflax, Dalmatian	Foliar herbicide treatment	No action	Foliar herbicide treatment	Remove flowering stalks
wild basil savory	Foliar herbicide treatment (most effective)	No action	No action	Foliar herbicide treatment (late summer)
willow-herb, hairy	Remove flowering stalks	Plants die back - no action	Foliar herbicide treatment	Remove flowering stalks
wormwood, absinth	No action	No action	Foliar herbicide treatment	Foliar herbicide treatment
yellow archangel	No action	No action	Foliar herbicide treatments	Foliar herbicide treatments
Category 2 Weeds				
bamboo, spp.	Foliar herbicide treatment	No action	Foliar herbicide treatment	No action
barberry, common	Cut stump herbicide treatment	No action	No action	Cut stump herbicide treatment

NOXIOUS WEED	FALL	WINTER	SPRING	SUMMER
bindweed, field	No action	Plants die back - no action	Foliar herbicide treatment	Foliar herbicide treatment
blackberry species	Foliar herbicide treatment on 1 year old stems only	Basal bark herbicide treatment	Foliar herbicide treatment; cut stump treatment (late spring)	No action
broom, Scotch	Cut stump or basal bark herbicide treatments	Cut stump or basal bark herbicide treatments	Cut stump or basal bark herbicide treatments; foliar treatments before flowering	Cut stump or basal bark herbicide treatments
burdock, common	Foliar herbicide treatment of rosettes; manual removal of flowering plants	No action	Foliar herbicide treatments	Manual removal of flowering plants
butterfly bush	Herbicide treatment - cut stump (large plants) or foliar (small plants)	No action	Herbicide treatment - cut stump (large plants) or foliar (small plants)	Herbicide treatment - cut stump (large plants) or foliar (small plants)
canarygrass, reed	Foliar herbicide treatment	Plants die back - no actions	Foliar herbicide treatment	No action
cheatgrass or downy brome	No action	No action	Foliar herbicide treatment	No action
chicory, common	Foliar herbicide treatment	No action	Foliar herbicide treatment	Foliar herbicide treatment
common mullein	No action	No action	Foliar herbicide treatment of rosettes	Manually remove bolting plants
hawthorn, English	Cut stump treatment (large plants) or foliar treatment (small plants)	No action	Cut stump treatment	Manual removal of small plants in early summer
herb Robert	Foliar herbicide treatment	No action	Foliar herbicide treatment	Foliar herbicide treatment
iris, yellow flag	Foliar herbicide treatment	Plants die back - no action	Foliar herbicide treatment	No action
Ivy, English	Cut stump or basal bark herbicide treatments	Cut stump or basal bark herbicide treatments	Manual removal when soil is moist; Cut stump or basal bark herbicide treatments	Cut stump or basal bark herbicide treatments

NOXIOUS WEED	FALL	WINTER	SPRING	SUMMER
mountain ash, European	Cut stump herbicide treatment	No action	Cut stump herbicide treatment	Cut stump herbicide treatment
mustard, wild	Foliar herbicide treatment of rosettes; Manual removal of flowering plants	Foliar herbicide treatment of rosettes	Foliar herbicide treatment of rosettes; Manual removal of flowering plants	No action – winter annual
Nightshade, hairy	Manual removal	No action – summer annual	Foliar herbicide treatment; manual removal	Manual removal
peavine, everlasting	Foliar herbicide treatments	Plants die back - no action	Foliar herbicide treatments	Foliar herbicide treatments
periwinkle, large	Foliar herbicide treatments	No action	Foliar herbicide treatments	Foliar herbicide treatments
St Johnswort, common	No action	No action	Foliar herbicide treatments	Foliar herbicide treatments
thistle, bull	Foliar herbicide treatment of rosettes	No action	Foliar herbicide treatment of rosettes	Foliar herbicide treatment of rosettes; manually remove flowering plants
thistle, Canada	Foliar herbicide treatment	Plants die back - no action	Foliar herbicide treatment	No action
toadflax, yellow	No action	No action	Foliar herbicide treatment	No action

2025 IWM TASKS

Biological
Coordinate with WSU Extension and Washington State Noxious Weed Control Board for studies on the effectiveness of biocontrols and participate in releases (if appropriate)
Physical
Update contact list to be shared between departments
Promote desirable native vegetation wherever possible
Posted Herbicide Application Notices to clearly mark treatment areas prior to all herbicide activity and leave up for at least 24 hours
Collaborate with Roads department, WSDOT and Clallam PUD to identify landscape goals and harmonize maintenance techniques
Re-imagine Adopt-A-Patch opportunities to increase public interest and participation
Review public involvement opportunities to ensure the available material meets program goals and is readily accessible online
Cultural
Further develop pollinator friendly plantings and coordinate with Road Department and WSU Extension to incorporate existing volunteer programs
Assist with research projects where possible
Continue building partnerships with local native plant sources and update material list to incorporate these projects
Partner with experts from local, state and federal agencies and entities including but not limited to: Clallam County Parks, Washington State University Extension, WSU Master Gardeners, local chapter of bee keepers, the native plant and Audubon societies, the Nature Conservancy, conservation districts, Olympic National Park, Olympic National Forest, USFW Marine Refuge System, Makah, Quileute, Lower Elwha Klallam, and Jamestown S'Klallam tribes, and others who have an interest in developing local native seed and plant resources for use in government projects
Manage "Owner Will Control" agreements. Maintain current list and map of "Owner Will Control" locations for both office and field use. Review "Owner Will Control" application process and forms to ensure all public involvement opportunities are readily accessible online. Encourage landowners with "Owner Will Control" agreements to undertake adjacent roadside enhancement consistent with developing a low maintenance, self-sustaining plant community to prevent weed invasion
Identify and compile a list of sites for revegetation opportunities
Develop native seed mix for Road Department projects where bare ground is necessary
Preventative
Update rock and gravel source weed management protocols
Continue our partnership with mowing personnel to facilitate practices that prevents the spread of propagules and encourages the growth of natural site appropriate vegetation

Identify County quarries, storage areas, and spoil disposal sites (pits) that will be actively used to store or process material and ensure stockpiles and adjacent areas are weed-free.
Survey roads that do not have up to date information on weeds located in right of way
Implement weed free material requirements for all county projects
Facilitate annual weed treatment overview and invasive plants identification training for Roads field crews
Identify equipment needs, investigate available resources, procure as resources allow
Treat all sand piles and sand extraction zones in county pits
Provide inspection services for all privately sourced material for county projects and provide the Roads Department with an up-to-date list of all private pits that meet the weed-free standard
Monitor and evaluate treatments in county pits
Catalog new weed infestations and update road priorities in treatment structure
Chemical
Implement project list based on tables 4-8
Post annual project list. Update during season as resources allow
Compile locations and instructions for special management areas. Include and update field maps as frequently as possible
Ready all necessary forms, regulatory compliance paperwork and safety equipment before commencing treatment season
Coordinate with Road Department staff to identify “special management areas” or non-native, invasive weed locations that interfere with road safety or function; outline additional management needs and strategy for weed control in these areas
Develop and utilize regional partners to assist in weed control across the county
Complete treatment records daily. Enter data into Clallam County Noxious Weed Control Program database. Monitor at least 10% of all treatments, retreat as needed and as resources allow
Identify any additional equipment needs and take steps to incorporate any available resources, including vehicles, application equipment, water tanks, or technical equipment
Provide WSU Master Gardeners Roadside Vegetation Monitoring Team (RVMT) with safety equipment, additional training opportunities, and technical support for monitoring projects
Conduct a weed inventory on at least 25% of all county roads annually
Identify, document and map regulated species, location, size and density
Update survey data of county roadsides and catalog infestations over time
Identify and compile a list of high priority infestations for the following year. Create maps if necessary.
Support, volunteer-based projects either on or adjoining county property that protects county property from weed infestations. This may include monitoring and revegetation projects
Promptly respond to all public inquiries. Address any public concerns regarding applications
Review and update on-line weed control request application process and forms as necessary
Review process for interdepartmental communication

Compile annual report summarizing accomplishments, effectiveness, and recommendations for subsequent year. Brief the Road Department and County Commissioners by December 31st

Draft IWM plan and submit to the Clallam County Noxious Weed Control Board and Road Department Supervisor for approval prior to the Weed Board's first meeting of the year. Submission of the IWM plan should occur 20 days before the meeting and should be posted online. Provide public notice that plan will be discussed, with weed board meeting announcements. The finalized plan and a map of proposed treatment locations should be posted online and made available upon public request

APPLICATION LOCATIONS

The 2025 Work Plan identifies treatment locations that may include the use of herbicides. The IWM Program priorities for roadside weed control in 2025 include:

- 1) Control of Category 1 regulated weeds on County roadsides in accordance with state law.
- 2) Control of Category 1 regulated weeds and select weeds at all County rock sources/District shops.
- 3) Control of Category 1 and 2 weeds at locations with most impact to local agriculture.
- 4) Control of Category 1 and 2 weeds at locations with most impact to local forestry.
- 5) Control of non-native weeds that interfere with the safety or function of County roadsides or additional non-roadside “special sites”.
- 6) Control of Category 1 and 2 weeds at locations requested by the public and local agencies.

In general, sites included in the 2025 Work Plan were chosen to complement previous work, to account for observed field conditions, or to further specific public benefit and need. Roadside treatments will include the most effective mix of methods as site or species appropriate. Tables 4-8 only include locations where herbicides may be used, possibly in combination with other treatment methods. The tables include county roads to their full extent; however, treatments may be limited to specific weed locations and in many cases only at isolated patches of Category 1 weeds.

Roads controlled in prior years for infestations of Category 1 weeds are included in this Work Plan for retreatment; adjacent roads may have also been included as needed. This Work Plan also includes county roads with isolated patches of Category 1 weed species as identified by past survey data. Isolated infestations can be controlled efficiently and require limited time and effort.

To prevent the spread of weeds, all county rock sources (pits/quarries), and district shops are included in the 2025 Plan as a priority for control.

This Work Plan includes control of roadside Category 2 weed targets in agriculture and forestry areas as identified in Tables 4-6; selections reflect input from farmers and foresters. Additional Category 2 weeds may be controlled as time and resources allow. Non-native, invasive weeds that interfere with the safety or function of the selected roads or weeds at additional County-owned “special sites” may be controlled as needed.

Early Detection Rapid Response (EDRR) sites are small roadside infestations, defined here as those that are patches that are less than 100 square feet per 1/10 of a mile, previously unknown, and/or requested by an affected owner or entity during the treatment season. The EDRR strategy is efficient, reduces costs, and increases efficacy of control. EDRR is a preferred strategy for this program, but by nature cannot be determined in advance. All sites that are treated under EDRR will be individually posted on site (with *Herbicide Notice*) and published online as soon as possible. Unknown county infestations found *outside* of the road right-of-way may be treated as EDRR sites up to 500 square feet.

The public may request control of Category 1 or Category 2 species at locations not listed in this plan. Control measures may include the use of herbicide, but more likely, will be controlled using the most effective mix of measures as site or species appropriate. Public requests will be accommodated with the highest priority given to Category 1 weeds, followed by Category 2 weeds in the order they are received. Requests received prior to March 1st are most likely to be accommodated in the same year as requested. Requests received after March 1st are likely to be verified and included in a subsequent year’s work plan.

SELECTED ROADS: TABLES 4-6

The Clallam County roadsides included in the 2025 Integrated Weed Management Plan are listed below in Tables 4 - 6. The tables are separated geographically into “East Clallam County”, “Central Clallam County” and “West Clallam County” and are intended to mirror the road maintenance district boundaries. Roads are grouped by similar weed management goals; a description of each “Group” is available at the beginning of the table.

Roads are assigned a relative “Treatment Priority” (1 - highest priority, 4 - lowest priority) based on infestation conditions and overall IWM program goals.

Priority 1 roads are intended to be completely treated in 2025

Priority 2 roads are intended to be treated in conjunction with adjacent roads or activities

Priority 3 roads will be treated as time and resources allow

Priority 4 roads have been put on a low maintenance cycle of 2-4 years for monitoring

EAST CLALLAM COUNTY SELECTED ROADS

East Clallam County includes areas around Carlsborg, Sequim, Blyn and the Miller Peninsula (reference Focus Area Maps 2-4). The Dungeness Valley, situated between Sequim and Carlsborg, is the primary agricultural region in Clallam County. Roadside infestations of Canada thistle impose significant impacts on adjacent agricultural producers and have been elevated in priority in East Clallam County to reflect the input and requests of local land managers.

Road Groups:

Agricultural: roadsides with weed infestations most likely to have an impact to adjacent agricultural producers; infestations of Canada thistle are intended to be completely treated.

Forestry: roadsides that have known weed infestations and are used to access forestry managed timber lands; increasing potential for cross contamination

Source Cat. 1 Weeds: roadsides with the most extensive infestations of category 1 weeds; treatment is intended to contain and reduce infestations.

Isolated Cat. 1 Weeds: roadsides with isolated infestations of category 1 weeds either treated or surveyed in the past; treatment is intended to eradicate individual infestations or plants.

Pit Entrance: roadsides leading to County pits; treatment is intended to contain and protect county resources.

Public Request: roadsides with infestations requested by individuals or public agencies; treatment is intended to meet requests as time and resources allow.

Survey: roadsides that should be surveyed for any new weeds every 2-4 years; this group also contains roads that have not been surveyed for noxious weeds

Table 4. East Clallam County roads selected for herbicide treatment in 2025

Target Weed Species For Control – additional non-native, invasive species may be controlled as time and resources allow. Species listed in *italics* are unconfirmed infestations.

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Blue Mountain Rd	Source Cat. 1 Weeds	7.0	meadow knapweed, tansy ragwort, bohemian knotweed	1
Business Park Loop	Source Cat. 1 Weeds	0.5	spotted knapweed	1
Carlsborg Rd	Agricultural	1.8	spotted knapweed, common tansy, spurge laurel	1
Cays Rd	Source Cat. 1 Weeds	2.8	Bohemian knotweed, Italian thistle, meadow knapweed, spotted knapweed, spurge laurel	1
Cline Spit Rd	Isolated Cat. 1 Weeds	0.3	poison hemlock	1
Coulter Rd	Isolated Cat. 1 Weeds	0.2	poison hemlock	1
Dawley Rd	Isolated Cat. 1 Weeds	0.7	poison hemlock, spurge laurel, tansy ragwort	1
Diamond Point Rd	Isolated Cat. 1 Weeds	4.0	poison hemlock, Scotch broom, tansy ragwort	1
Eagle Creek Rd-West 400ft	Isolated Cat. 1 Weeds	0.1	tansy ragwort	1
East Anderson Rd	Agricultural	2.1	poison hemlock, common tansy	1
East Sequim Bay Rd	Isolated Cat. 1 Weeds	4.5	wild basil savory, tansy ragwort	1
Evans Rd	Agricultural	1.7	common teasel, poison hemlock	1
Fasola Rd	Isolated Cat. 1 Weeds	1.0	common teasel, poison hemlock, common tansy	1
Gardiner Beach Rd-West 400ft	Survey	0.1	No target weeds found	1
Happy Valley Rd	Source Cat. 1 Weeds	5.7	common teasel, meadow knapweed, poison hemlock, tansy ragwort	1
Johnson Creek Rd	Forestry	0.5	meadow knapweed	1
Kitchen-Dick Rd	Agricultural	3.2	absinth wormwood, common tansy, common teasel, hairy whitetop, hoary alyssum, meadow knapweed, poison hemlock, spotted knapweed, tansy ragwort, spurge laurel	1
Lamar Ln	Source Cat. 1 Weeds	0.7	Italian thistle, poison hemlock	1
Lost Mountain Rd	Forestry	5.0	meadow knapweed, tansy ragwort, common teasel	1
Lotzgesell Rd	Agricultural	3.4	common tansy, common teasel, meadow knapweed, spotted knapweed, poison hemlock	1
Marine Dr	Isolated Cat. 1 Weeds	1.6	poison hemlock	1
Marmot Loop	Isolated Cat. 1 Weeds	0.5	wild chervil	1
North Brown Rd -North of East Hendrickson Rd	Isolated Cat. 1 Weeds	0.5	common teasel, meadow knapweed, tansy ragwort	1
Old Blyn Hwy	Isolated Cat. 1 Weeds	2.1	poison hemlock, tansy ragwort, wild basil savory, common tansy	1
Palo Alto Rd	Forestry	8.4	meadow knapweed, sulfur cinquefoil, tansy ragwort, wild basil savory, poison hemlock	1
Port Williams Rd	Agricultural	2.3	common teasel, tansy ragwort, poison hemlock	1

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
River Rd	Source Cat. 1 Weeds	2.5	common fennel, common tansy, meadow knapweed, spotted knapweed, sulfur cinquefoil	1
Sequim-Dungeness Way (North of 1323 address)	Agricultural	4.1	common fennel, common teasel, meadow knapweed, poison hemlock, spotted knapweed, tansy ragwort, common tansy	1
South 3rd Ave (Reservoir Rd to Happy Valley Rd)	Isolated Cat. 1 Weeds	0.5	meadow knapweed, mole plant, tansy ragwort	1
Towne Rd	Agricultural	3.0	common fennel, common teasel, hairy whitetop, meadow knapweed, spotted knapweed, poison hemlock	1
Turnstone Ln	Source Cat. 1 Weeds	0.7	spotted knapweed, meadow knapweed	1
Vistas Dr	Pit Entrance	0.2	poison hemlock, Scotch broom	1
Ward Rd	Agricultural	1.7	Bohemian knotweed, poison hemlock, yellow archangel	1
Washington Harbor Rd	Isolated Cat. 1 Weeds	1.2	Bohemian knotweed, poison hemlock,	1
West Washington St	Source Cat. 1 Weeds	0.3	poison hemlock, spotted knapweed, spurge laurel	1
Woods Rd	Forestry	1.7	tansy ragwort, wild basil savory, common tansy, herb Robert	1
3 Crabs Rd	Isolated Cat. 1 Weeds	1.4	poison hemlock, common fennel, common tansy	2
Aster Rd	Isolated Cat. 1 Weeds	0.1	tansy ragwort	2
Atterberry Rd	Isolated Cat. 1 Weeds/Public request	2.8	meadow knapweed, poison hemlock, tansy ragwort, Scotch broom	2
Barker Rd	Isolated Cat. 1 Weeds	0.1	poison hemlock	2
Born Dr	Isolated Cat. 1 Weeds	0.1	common tansy	2
Brittany Ln	Survey	0.3	No target weeds found	2
Buckhorn Rd	Isolated Cat. 1 Weeds	0.4	hairy whitetop	2
Cameron Rd-District 1	Agricultural	0.8	poison hemlock	2
Cassidy Rd	Forestry	0.8	poison hemlock, spotted knapweed, spurge laurel, tansy ragwort, meadow knapweed	2
Cat Lake Rd	Public Request	1.2	poison hemlock, Scotch broom, tansy ragwort	2
Chicken Coop Rd	Forestry	3.5	common teasel, tansy ragwort, wild basil savory	2
Cormorant Dr	Survey	0.1	No target weeds found	2
Corriea Rd	Isolated Cat. 1 Weeds	1.0	common teasel, poison hemlock	2
Dickinson St	Isolated Cat. 1 Weeds	0.2	poison hemlock	2
Discovery View Dr	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Don Schmith Rd	Isolated Cat. 1 Weeds	0.5	meadow knapweed, sulfur cinquefoil, tansy ragwort	2
East Nelson Rd	Survey	0.3	No target weeds found	2
East Runnion Rd	Public request	0.8	Scotch broom, spotted knapweed	2
East Silberhorn Rd (Viking Wy to S 7 th Ave)	Survey	0.6	No target weeds found	2

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Easterly Rd	Isolated Cat. 1 Weeds	0.3	meadow knapweed, tansy ragwort	2
Eggloff Rd	Forestry	0.4	meadow knapweed, tansy ragwort	2
Finn Hall Rd	Isolated Cat. 1 Weeds	2.5	common teasel, poison hemlock	2
Fish Hatchery Rd	Forestry	1.8	Bohemian knotweed, yellow archangel	2
Fleming Dr	Isolated Cat. 1 Weeds	0.4	poison hemlock	2
Gasman Rd	Public request	1.5	tansy ragwort, Scotch broom	2
Gellor Rd	Isolated Cat. 1 Weeds	1.0	meadow knapweed	2
Gilbert Rd	Isolated Cat. 1 Weeds	0.2	poison hemlock, spurge laurel	2
Greywolf Rd	Survey	0.5	No target weeds found	2
Gunn Rd	Agricultural	0.9	No target weeds found	2
Gupster Rd	Isolated Cat. 1 Weeds	0.5	poison hemlock, spotted knapweed, meadow knapweed, spurge laurel	2
Heath Rd	Survey	1.5	No target weeds found	2
Hogback Rd	Agricultural	0.8	Canada thistle, Scotch broom	2
Holgerson Rd	Isolated Cat. 1 Weeds	0.4	common teasel	2
Holland Rd	Agricultural	1.7	yellow archangel	2
Hooker Rd	Isolated Cat. 1 Weeds	2.7	poison hemlock, common teasel	2
Industrial Pkwy	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Jamestown Rd	Isolated Cat. 1 Weeds	1.6	common fennel	2
Jimmy Come Lately Rd	Forestry	1.2	meadow knapweed, Bohemian knotweed, herb Robert	2
Kirk Rd	Survey	0.3	No target weeds found	2
Kirner Rd	Pit Entrance	0.9	meadow knapweed, common tansy	2
Louella Rd	Isolated Cat. 1 Weeds	0.9	meadow knapweed, tansy ragwort	2
Lupine Dr	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Madrona Way	Isolated Cat. 1 Weeds	1.1	tansy ragwort	2
Manzanita Dr	Isolated Cat. 1 Weeds	0.7	tansy ragwort	2
Matson Rd	Agricultural	0.5	No target weeds found	2
North St	Isolated Cat. 1 Weeds	0.4	tansy ragwort, spurge laurel	2
North Priest St	Isolated Cat. 1 Weeds	0.3	meadow knapweed, poison hemlock	2
Old Olympic Hwy	Source Cat. 1 Weeds	9.6	absinth wormwood, common tansy, meadow knapweed, poison hemlock, spotted knapweed, spurge laurel, tansy ragwort, common teasel, moleplant	2
Olympic Vista	Isolated Cat. 1 Weeds	0.1	spurge laurel	2
Panorama Blvd	Source Cat. 1 Weeds	0.8	tansy ragwort	2
Parrish Rd	Survey	1.0	No target weeds found	2
Pierce Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	2
Pinnell Rd	Public request	0.7	Canada thistle, meadow knapweed	2
Poplar Ct	Isolated Cat. 1 Weeds	0.1	tansy ragwort	2

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Rhododendron Dr	Isolated Cat. 1 Weeds	0.8	poison hemlock, tansy ragwort	2
Ridge View Dr	Isolated Cat. 1 Weeds	0.8	poison hemlock	2
Salal Way	Isolated Cat. 1 Weeds	0.4	tansy ragwort	2
Sawmill Rd	Survey	0.3	N/A	2
Schindler Rd	Isolated Cat. 1 Weeds	0.3	poison hemlock, tansy ragwort	2
Schmuck Rd	Agricultural	1.3	common teasel	2
Schoolhouse Point Ln	Isolated Cat. 1 Weeds	0.4	tansy ragwort	2
Senz Rd	Isolated Cat. 1 Weeds	0.5	spurge laurel	2
Sherwood Rd	Isolated Cat. 1 Weeds	0.3	poison hemlock, tansy ragwort	2
Siebert Creek Rd	Isolated Cat. 1 Weeds	0.8	sulfur cinquefoil	2
Smithfield Dr	Isolated Cat. 1 Weeds	0.3	spotted knapweed	2
Sophus Rd	Survey	0.1	No target weeds found	2
South St	Public Request	0.1	Scotch broom	2
Sturdevant Rd	Survey	0.7	No target weeds found	2
Sunshine Ave	Isolated Cat. 1 Weeds	0.6	tansy ragwort	2
Sunshine Plz	Isolated Cat. 1 Weeds	0.2	tansy ragwort	2
Taylor Cut-off Rd	Isolated Cat. 1 Weeds	2.7	meadow knapweed, spotted knapweed, spurge laurel	2
Taylor Ranch Rd	Isolated Cat. 1 Weeds	0.8	Italian arum, tansy ragwort, poison hemlock, meadow knapweed	2
Thompson Rd-District 1	Forestry	1.8	poison hemlock, Scotch broom, tansy ragwort, common tansy	2
Thornton Dr	Isolated Cat. 1 Weeds	1.5	poison hemlock, tansy ragwort	2
Timber Rd	Isolated Cat. 1 Weeds	0.1	tansy ragwort	2
Tripp Rd	Isolated Cat. 1 Weeds	0.3	orange hawkweed	2
Twin View Dr	Isolated Cat. 1 Weeds	0.6	meadow knapweed	2
Vista View Dr	Isolated Cat. 1 Weeds	0.3	poison hemlock, yellow archangel	2
West Anderson Rd	Survey	0.8	No target weeds found	2
West Hendrickson Rd	Agricultural	1.0	common teasel, poison hemlock	2
West Johnson Dr	Survey	0.1	N/A	2
West Nelson Rd	Survey	0.6	No target weeds found	2
West Sequim Bay Rd	Isolated Cat. 1 Weeds	0.9	spotted knapweed, tansy ragwort, common teasel, poison hemlock, spurge laurel	2
Wilcox Ln	Survey	0.7	N/A	2
Woodcock Rd	Isolated Cat. 1 Weeds	6.3	poison hemlock, hairy whitetop, spurge laurel, common teasel, common fennel	2
Woodland Dr	Isolated Cat. 1 Weeds	0.4	spurge laurel	2

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Youngquist Rd	Isolated Cat. 1 Weeds	0.9	tansy ragwort, spurge laurel, common teasel	2
Zaccardo Rd	Isolated Cat. 1 Weeds	0.4	poison hemlock, tansy ragwort	2
Alderwood Dr	Survey	0.2	No target weeds found	3
Annabelle Ln	Survey	0.4	No target weeds found	3
Autumn Rd	Isolated Cat. 1 Weeds	0.6	spotted knapweed	3
Barbara St	Survey	0.3	No target weeds found	3
Barnes Rd	Survey	0.7	No target weeds found	3
Bellway Rd	Survey	0.2	No target weeds found	3
Buck Loop Rd	Isolated Cat. 1 Weeds	0.5	tansy ragwort	3
Cassidy Creek Rd	Survey	0.9	No target weeds found	3
Charles Way	Survey	0.1	No target weeds found	3
Clark Rd	Agricultural	0.9	poison hemlock, tansy ragwort	3
Cline Rd	Survey	0.6	No target weeds found	3
Conner Rd	Survey	0.3	No target weeds found	3
Crestline Dr	Survey	0.1	No target weeds found	3
Dryke Rd	Survey	0.7	No target weeds found	3
East Belfield Ave	Survey	0.2	No target weeds found	3
East Blair Ave	Isolated Cat. 1 Weeds	0.3	spurge laurel	3
East Bluff Dr	Isolated Cat. 1 Weeds	0.6	tansy ragwort	3
Elliott Ct	Survey	0.1	N/A	3
Emery Rd	Survey	0.6	No target weeds found	3
Forest Ridge Dr	Survey	0.2	No target weeds found	3
Frick St	Survey	0.1	No target weeds found	3
Frost Rd	Survey	0.5	No target weeds found	3
Gehrke Rd	Survey	0.9	No target weeds found	3
Grandview Dr	Survey	0.5	No target weeds found	3
Guiles Rd	Survey	0.5	N/A	3
Hardwick Rd	Isolated Cat. 1 Weeds	0.4	tansy ragwort	3
Heuhslein Rd	Survey	1.1	No target weeds found	3
Island Vista Way	Survey	0.3	No target weeds found	3
John Scott Rd	Isolated Cat. 1 Weeds	0.2	poison hemlock	3
Joslin Rd	Survey	0.4	No target weeds found	3
Juan de Fuca Way	Survey	0.2	No target weeds found	3
Knapp Rd	Survey	0.2	No target weeds found	3
Lilly Rd	Survey	0.2	No target weeds found	3
Lester Way	Survey	0.2	No target weeds found	3

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Lewis Rd	Survey	0.6	No target weeds found	3
Lochow Rd	Survey	0.2	No target weeds found	3
Madrona Ter	Survey	0.3	No target weeds found	3
Mariott Ave (MP 0.08 to 0.3)	Survey	0.2	N/A	3
Marshall Rd	Survey	0.5	No target weeds found	3
McAlmond St	Survey	0.1	No target weeds found	3
McComb Rd	Agricultural	0.8	common teasel	3
McFarland Dr	Survey	1.2	No target weeds found	3
Medsker Rd	Survey	1.3	No target weeds found	3
Mill Rd	Survey	0.6	No target weeds found	3
Montrose Pl	Survey	0.1	No target weeds found	3
Mountain View Dr	Survey	0.4	No target weeds found	3
North Barr Rd	Isolated Cat. 1 Weeds	1.3	tansy ragwort, poison hemlock	3
North Boyce Rd	Survey	0.1	No target weeds found	3
North Eldridge Rd	Isolated Cat. 1 Weeds	0.2	tansy ragwort	3
North Kendall Rd	Survey	0.4	No target weeds found	3
Olson Rd	Survey	1.8	No target weeds found	3
Olstead Rd	Survey	0.5	No target weeds found	3
Peninsula St	Survey	0.1	No target weeds found	3
Riverside Rd	Isolated Cat. 1 Weeds	0.5	poison hemlock	3
Sandstone Pl	Survey	0.1	No target weeds found	3
Secor Rd	Isolated Cat. 1 Weeds	0.5	spotted knapweed	3
Serpentine Ave	Isolated Cat. 1 Weeds	0.8	poison hemlock	3
Sherburne Rd	Isolated Cat. 1 Weeds	1.0	spotted knapweed, sulfur cinquefoil, spurge laurel	3
Shore Rd	Survey	0.9	No target weeds found	3
South Barr Rd	Isolated Cat. 1 Weeds	0.7	poison hemlock	3
South Boyce Rd	Survey	0.2	No target weeds found	3
South Eldridge Rd	Survey	0.4	No target weeds found	3
Spath Rd	Public request	1.3	Himalayan blackberry	3
Spencer Rd	Survey	0.25	No target weeds found	3
Sporseen Rd	Isolated Cat. 1 Weeds	0.9	common teasel	3
Spring Rd	Survey	0.6	No target weeds found	3
Stuart Dr	Survey	0.4	No target weeds found	3
Summerset Ct	Survey	0.1	N/A	3
Sylvan Way	Survey	0.2	No target weeds found	3
Vautier Rd	Agricultural	0.6	No target weeds found	3

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Voice Of America Rd	Isolated Cat. 1 Weeds	1.3	poison hemlock	3
Waldo Rd	Survey	0.2	No target weeds found	3
West Alder Ct	Survey	0.1	No target weeds found	3
West Bluff Dr	Survey	0.3	No target weeds found	3
West Palo Verde Loop	Survey	0.3	No target weeds found	3
West Runnion Rd	Isolated Cat. 1 Weeds	0.5	poison hemlock, spotted knapweed	3
West Silberhorn Rd	Isolated Cat.1 Weeds	0.5	poison hemlock, common fennel	3
West Spruce Ct (N Priest Rd to 1000 W Spruce Ct)	Survey	0.3	No target weeds found	3
Wild Currant Way	Survey	0.3	No target weeds found	3
Abbott Rd	Survey	0.2	No target weeds found	4
Applegate Ln	Survey	0.3	No target weeds found	4
Balmoral Ct	Survey	0.1	No target weeds found	4
Bay View St	Survey	0.4	No target weeds found	4
Brigadoon Blvd	Survey	0.9	No target weeds found	4
Christian Ln	Survey	0.2	No target weeds found	4
Cook Rd	Survey	0.3	No target weeds found	4
Cottonwood Ln	Survey	0.2	No target weeds found	4
Dent Pl	Survey	0.1	N/A	4
Deytona St	Survey	0.3	N/A	4
Eagle Pl	Survey	0.1	No target weeds found	4
East Bedinger Rd	Survey	0.1	No target weeds found	4
East Diane Rd	Survey	0.1	No target weeds found	4
East Robert Pl	Survey	0.1	No target weeds found	4
Eberle Ln	Survey	0.3	No target weeds found	4
Elizabeth Ln	Survey	0.4	No target weeds found	4
Farm Pl	Survey	0.1	No target weeds found	4
Fitzgerald Rd	Survey	0.2	No target weeds found	4
Garden Ln	Survey	0.1	No target weeds found	4
House Rd	Survey	0.4	No target weeds found	4
Hudon Rd	Survey	0.2	No target weeds found	4
Jake Hall Rd	Survey	0.4	No target weeds found	4
James St	Survey	0.1	No target weeds found	4
Jay Robinson Rd	Survey	0.2	No target needs found	4
Josephine Pl	Survey	0.1	No target weeds found	4

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
June Pl	Survey	0.1	No target weeds found	4
Karen Ct	Survey	0.1	No target weeds found	4
Karie Ct	Survey	0.1	No target weeds found	4
Kellie Ct	Survey	0.1	No target weeds found	4
Koeppe Dr	Survey	0.2	No target weeds found	4
Lange Ln	Survey	0.3	No target weeds found	4
Libby St	Survey	0.4	No target weeds found	4
Lighthouse View Dr	Survey	0.1	No target weeds found	4
Lorraine Dr	Survey	0.1	No target weeds found	4
Mains Rd	Survey	0.3	No target weeds found	4
Mantle Rd	Survey	0.4	No target weeds found	4
Marchbanks Rd	Survey	0.1	No target weeds found	4
Macleay Rd	Survey	0.6	No target weeds found	4
Meadow Lark Ln	Survey	0.3	No target weeds found	4
Moonlight Dr	Survey	0.4	No target weeds found	4
Nello Pl	Survey	0.1	No target weeds found	4
Nicole Pl	Survey	0.2	No target weeds found	4
North Olympic View Ave	Survey	0.2	No target weeds found	4
North Scott Dr	Survey	0.1	No target weeds found	4
Park Ln	Survey	0.1	No target weeds found	4
Percy Ln	Survey	0.1	No target weeds found	4
Pebble Way	Survey	0.1	No target weeds found	4
Phinn Rd	Survey	0.1	No target weeds found	4
Robbins Rd	Survey	0.3	No target weeds found	4
Ridge Pl	Survey	0.1	No target weeds found	4
Schmith Rd	Survey	0.2	No target weeds found	4
Simpson Rd	Survey	0.4	No target weeds found	4
Sofie Rd	Survey	0.2	No target weeds found	4
South Olympic View Ave	Survey	0.4	No target weeds found	4
South Scott Dr	Survey	0.4	No target weeds found	4
Starry Rd	Survey	0.2	No target weeds found	4
Steve Pl	Survey	0.1	No target weeds found	4
Stone Rd	Survey	0.6	No target weeds found	4
Streit Rd	Survey	0.2	No target weeds found	4
Sundial Loop	Survey	0.3	No target weeds found	4
Tyee Rd	Survey	0.2	No target weeds found	4
Van Dyken Dr	Survey	0.3	No target weeds found	4

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
West Bedinger Rd	Survey	0.1	No target weeds found	4
West Diane Dr	Survey	0.1	No target weeds found	4
West Oak Ct	Survey	0.3	No target weeds found	4
West Robert Pl	Survey	0.1	No target weeds found	4
Wheeler Rd	Survey	0.5	No target weeds found	4
Willard Dr	Survey	0.1	No target weeds found	4
Williamson Rd	Survey	0.6	No target weeds found	4
Wright Rd	Survey	0.5	No target weeds found	4
Total: 273 roads		Total: 213.65 mi		

¹Target Weed Species for Control – additional non-native, invasive species may be controlled as time and resources allow.

CENTRAL CLALLAM COUNTY SELECTED ROADS

Central Clallam County includes the approximate areas of Port Angeles, Joyce and the Elwha Valley (reference Focus Area Maps 5-6). The roads selected for treatment in 2025 primarily include infestations of Category 1 weed species and adjacent roads. Significant roadside infestations of meadow knapweed, knotweed and tansy ragwort have been previously treated and require retreatment and monitoring in 2025.

Road Groups:

Clallam Co Restoration Project: roadsides adjacent to Clallam County environmental restoration or mitigation projects and may include native revegetation efforts; treatment is intended to contain infestations of non-native, invasive species to protect native plantings and function.

Source Cat. 1 Weeds: roadsides with the most extensive infestations of category 1 weeds; treatment is intended to contain and reduce infestations.

Isolated Cat. 1 Weeds: roadsides with isolated infestations of category 1 weeds either treated or surveyed in the past; treatment is intended to eradicate individual infestations or plants.

Forestry: roadsides that have known weed infestations and are used to access forestry managed timber lands; increasing potential for cross contamination

Pit Entrance: roadsides leading to County pits; treatment is intended to contain and protect county resources.

Public Request: roadsides with infestations requested by individuals or public agencies; treatment is intended to meet requests as time and resources allow.

Survey: roadsides that should be surveyed for any new weeds every 2-4 years; this group also contains roads that have not been surveyed for noxious weeds

Table 5. Central Clallam County roads selected for herbicide treatment in 2025

Target Weed Species For Control – additional non-native, invasive species may be controlled as time and resources allow. Species listed in *italics* are unconfirmed infestations.

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Benson Rd	Isolated Cat. 1 Weeds	1.1	orange hawkweed	1
Black Diamond Rd	Forestry	4.4	common teasel, meadow knapweed, wild basil savory, common tansy	1
Camp Hayden Rd	Forestry	3.5	tansy ragwort, meadow knapweed, wild basil savory	1
Crescent Beach Rd	Forestry	3.5	tansy ragwort, meadow knapweed	1
Dan Kelly Rd	Forestry	3.2	Bohemian knotweed, tansy ragwort, meadow knapweed, common tansy, yellow archangel	1
Deer Park Loop	Clallam Co Restoration Project	0.6	common teasel, meadow knapweed, Scotch broom	1
East Beach Rd	Source Cat. 1 Weeds	0.8	meadow knapweed, tansy ragwort, Scotch broom, spurge laurel	1

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
East Lyre River Rd	Isolated Cat. 1 Weeds	0.6	meadow knapweed, tansy ragwort, wild basil savory	1
Eden Valley Rd	Isolated Cat. 1 Weeds	3.5	Bohemian knotweed, common teasel, meadow knapweed, tansy ragwort	1
Elwha River Rd	Forestry	1.9	Scotch broom, meadow knapweed, tansy ragwort, herb Robert	1
Farrington Rd	Source Cat. 1 Weeds	0.9	meadow knapweed, tansy ragwort	1
Fisher Cove Rd	Forestry	0.8	Bohemian knotweed, common tansy, meadow knapweed, wild basil savory	1
Freshwater Bay Rd	Isolated Cat. 1 Weeds	2.5	tansy ragwort	1
Freshwater Park	Isolated Cat. 1 Weeds	1.3	tansy ragwort	1
Gagnon Rd	Isolated Cat. 1 Weeds	0.5	Bohemian knotweed, meadow knapweed, tansy ragwort	1
Granite Rd	Isolated Cat. 1 Weeds	0.3	meadow knapweed	1
Gravel Pit Rd	Pit Entrance	0.1	Scotch broom, common tansy, herb Robert	1
Herrick Rd	Pit Entrance	1.5	meadow knapweed, herb Robert, Scotch broom, Bohemian knotweed, spurge laurel	1
Joyce-Piedmont Rd	Forestry	4.0	Bohemian knotweed, meadow knapweed, tansy ragwort, yellow archangel, common tansy, herb Robert	1
Laird Rd	Isolated Cat. 1 Weeds	0.9	meadow knapweed, tansy ragwort, Himalayan blackberry	1
Lake Dawn Rd	Isolated Cat. 1 Weeds	0.5	orange hawkweed, spurge laurel	1
Little Loop Dr	Isolated Cat. 1 Weeds	0.8	tansy ragwort	1
Little River Rd	Forestry	7.2	common teasel, meadow knapweed, tansy ragwort, yellow archangel, wild basil savory	1
Lower Elwha Rd	Clallam Co Restoration Project	2.8	meadow knapweed, Scotch broom, tansy ragwort	1
McGarvie Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	1
Mount Pleasant Rd	Forestry	5.7	meadow knapweed, poison hemlock, tansy ragwort, common tansy, mountain knapweed	1
O'Brien Rd	Isolated Cat. 1 Weeds	4.1	meadow knapweed, tansy ragwort, common tansy, sulfur cinquefoil	1
Okerman Rd	Isolated Cat. 1 Weeds	0.3	meadow knapweed	1
Olympic Hot Springs Rd	Isolated Cat. 1 Weeds	1.0	Bohemian knotweed, meadow knapweed, tansy ragwort, wild basil savory	1
Park and Ride Rd	Clallam Co Restoration Project	0.1	Himalayan blackberry, Scotch broom, tree lupine	1
Place Rd	Isolated Cat. 1 Weeds	2.5	Scotch broom, tansy ragwort	1
Schmitt Rd	Isolated Cat. 1 Weeds	0.5	tansy ragwort	1
South Shore Rd	Isolated Cat. 1 Weeds	2.1	wild basil savory, meadow knapweed, orange hawkweed	1
Township Line Rd	Forestry	1.7	tansy ragwort, meadow knapweed, yellow hawkweed, Bohemian knotweed	1
West Edgewood Dr	Isolated Cat. 1 Weeds	2.5	common tansy, meadow knapweed, tansy ragwort, common fennel, yellow archangel	1
West Lyre River Rd	Isolated Cat. 1 Weeds	0.6	meadow knapweed, tansy ragwort	1

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Whiskey Creek Beach Rd (Hwy 112 to Schmitt Rd)	Isolated Cat. 1 Weeds	0.5	tansy ragwort	1
Airport Rd	Isolated Cat. 1 Weeds	0.7	common teasel, meadow knapweed, tansy ragwort	2
Alice Rd	Survey	0.5	No target weeds found	2
Baker Farm Rd	Survey	0.4	No target weeds found	2
Bean Rd	Isolated Cat. 1 Weeds	0.5	meadow knapweed	2
Billy Smith Rd	Survey	0.7	No target weeds found	2
Bishop Rd	Isolated Cat. 1 Weeds	0.4	tansy ragwort, Himalayan blackberry, meadow knapweed	2
Cameron Rd -District 2	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Deer Park Rd	Forestry	8.7	tansy ragwort, yellow hawkweed, spurge laurel, meadow knapweed, common fennel	2
Dempsey Rd	Isolated Cat. 1 Weeds	0.7	tansy ragwort	2
Dietz Rd	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Dry Creek Rd	Isolated Cat. 1 Weeds	0.7	meadow knapweed, tansy ragwort	2
Durrwachter Rd	Isolated Cat. 1 Weeds	1.0	tansy ragwort	2
East Ennis Creek Rd	Isolated Cat. 1 Weeds	0.5	common tansy, yellow archangel	2
East Scrivner Rd	Isolated Cat. 1 Weeds	1.1	Bohemian knotweed	2
Evergreen View Pkwy	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Fey Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	2
Frederickson Rd	Survey	0.3	No target weeds found	2
Gerber Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	2
Glass Rd	Isolated Cat. 1 Weeds	1.2	Bohemian knotweed, tansy ragwort	2
Gossett Rd	Isolated Cat. 1 Weeds	0.9	meadow knapweed, tansy ragwort, yellow archangel	2
Harry Brown Rd	Isolated Cat. 1 Weeds	0.2	tansy ragwort	2
Hauk Rd	Isolated Cat. 1 Weeds	0.2	Bohemian knotweed	2
Henry Boyd Rd	Isolated Cat. 1 Weeds	0.7	Bohemian knotweed, tansy ragwort	2
Hoare Rd	Isolated Cat. 1 Weeds	0.7	meadow knapweed	2
Huckleberry Hill Dr	Isolated Cat. 1 Weeds	0.4	tansy ragwort	2
James Page Rd	Public Request	0.3	Canada thistle	2
Kacee Way	Isolated Cat. 1 Weeds	0.4	meadow knapweed	2
Karpen Rd	Survey	0.7	No target weeds found	2
Kemp St	Isolated Cat. 1 Weeds	0.8	Bohemian knotweed	2
Kreaman Rd	Survey	0.3	No target weeds found	2

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Lake Aldwell Rd	Isolated Cat. 1 Weeds	0.6	meadow knapweed, herb Robert	2
Liljedahl Rd	Isolated Cat. 1 Weeds	1.0	tansy ragwort	2
Mapleton Way	Isolated Cat. 1 Weeds	0.3	tansy ragwort	2
McCarver St	Survey	0.5	No target weeds found	2
Monroe Rd	Isolated Cat. 1 Weeds	2.9	tansy ragwort, meadow knapweed, common tansy	2
Mount Angeles Rd (South of Eckard Rd)	Forestry	4.4	tansy ragwort, meadow knapweed, Bohemian knotweed, orange hawkweed	2
Myrtle St	Isolated Cat. 1 Weeds	0.3	common tansy	2
North Bay View Ave	Isolated Cat. 1 Weeds	0.1	common tansy, Scotch broom	2
North Brook Ave	Public Request	0.6	Himalayan blackberry	2
North Masters Rd	Public request	0.5	Scotch broom	2
Old Black Diamond	Isolated Cat. 1 Weeds	0.2	meadow knapweed, tansy ragwort	2
Old Mill Rd (South of 4704 address)	Forestry	1.0	Bohemian knotweed, meadow knapweed	2
Oxenford Rd	Survey	1.0	No target weeds found	2
Power Plant Rd	Isolated Cat. 1 Weeds	0.8	Bohemian knotweed, meadow knapweed, tansy ragwort, Himalayan blackberry	2
Ranger Rd	Survey	1.0	No target weeds found	2
Rife Rd	Isolated Cat. 1 Weeds	0.6	Bohemian knotweed	2
Simmons Rd	Survey	1.0	No target weeds found	2
South Aviation Pl	Isolated Cat. 1 Weeds	0.1	meadow knapweed	2
South Bagley Creek Rd	Isolated Cat. 1 Weeds	1.5	Bohemian knotweed, poison hemlock	2
South Tiller Rd	Isolated Cat. 1 Weeds	0.2	Bohemian knotweed	2
Striped Peak Rd	Forestry	0.2	tansy ragwort	2
Wasankari Rd	Isolated Cat. 1 Weeds	0.6	tansy ragwort	2
West Arnette Rd	Isolated Cat. 1 Weeds	0.1	common teasel, Bohemian knotweed	2
West Duval Pl	Isolated Cat. 1 Weeds	0.2	tansy ragwort, meadow knapweed	2
Wye Rd	Isolated Cat. 1 Weeds	0.6	tansy ragwort	2
Avis St	Survey	0.4	No target weeds found	3
Bay St	Survey	0.6	No target weeds found	3
Brown Rd	Isolated Cat. 1 Weeds	0.6	meadow knapweed	3
Buchanan Dr	Survey	0.7	No target weeds found	3
Cedar Park Dr	Isolated Cat. 1 Weeds	0.7	tansy ragwort	3

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Champion Rd	Survey	0.2	No target weeds found	3
Chinook Ln	Survey	0.1	No target weeds found	3
Critchfield Rd	Survey	0.3	No target weeds found	3
Crown Z Water Rd	Public Request	0.4	Scotch broom	3
Doss Rd	Isolated Cat. 1 Weeds	0.5	tansy ragwort	3
Draper Rd	Isolated Cat. 1 Weeds	1.0	tansy ragwort, poison hemlock	3
Dunmire Rd	Survey	0.3	No target weeds found	3
East Arnette Rd	Survey	0.4	No target weeds found	3
Elwha Bluffs Rd	Survey	0.3	No target weeds found	3
Erving Jacobs Rd	Survey	0.5	No target weeds found	3
Fors Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	3
Grael-Ramapo Rd	Isolated Cat. 1 Weeds	1.5	tansy ragwort	3
Harrington Rd	Survey	0.2	No target weeds found	3
Hidden Valley Rd	Survey	0.5	No target weeds found	3
Hoffman Rd	Isolated Cat. 1 Weeds	0.2	tansy ragwort	3
Howard Rd	Isolated Cat. 1 Weeds	0.6	tansy ragwort	3
Hulse Rd	Survey	0.1	No target weeds found	3
John Jacobs Rd	Isolated Cat. 1 Weeds	0.3	poison hemlock	3
Key Rd	Survey	0.6	No target weeds found	3
King St	Survey	0.3	No target weeds found	3
Lake Farm Rd	Survey	0.7	No target weeds found	3
Leighland Ave	Survey	0.5	No target weeds found	3
Lawrence Rd	Isolated Cat. 1 Weeds	0.8	tansy ragwort	3
Marsden Rd	Survey	0.5	No target weeds found	3
Melton Rd	Isolated Cat. 1 Weeds	0.3	orange hawkweed	3
Miles Rd	Survey	0.3	No target weeds found	3
Miller Rd	Survey	0.7	No target weeds found	3
North Evergreen Dr	Survey	0.5	No target weeds found	3
North Larch Ave	Survey	0.5	No target weeds found	3
North Lee's Creek Rd	Survey	0.6	No target weeds found	3
Patterson Rd	Survey	1.0	No target weeds found	3
Pearce Rd	Isolated Cat, 1 Weeds	0.7	tansy ragwort	3
Peters Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	3
Reddick Rd	Survey	0.6	No target weeds found	3

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Reservoir Rd	Isolated Cat. 1 Weeds	0.2	tansy ragwort	3
Reynolds Rd	Isolated Cat. 1 Weeds	0.4	meadow knapweed, tansy ragwort	3
Robinson Rd	Survey	0.4	No target weeds found	3
Pavel Rd	Survey	0.5	No target weeds found	3
Shadow Ln	Isolated Cat. 1 Weeds	0.4	Bohemian knotweed, tansy ragwort	3
Silver Ln	Survey	0.1	No target weeds found	3
Sisson Rd	Survey	0.3	No target weeds found	3
Slate Dr	Survey	0.1	No target weeds found	3
South Bay View Ave	Survey	0.3	No target weeds found	3
South Brook Ave	Survey	0.2	No target weeds found	3
South Evergreen Dr	Isolated Cat. 1 Weeds	0.1	tansy ragwort	3
Sutter Rd	Survey	0.6	No target weeds found	3
Tamarack Ln	Isolated Cat. 1 Weeds	0.6	yellow archangel	3
Thompson Rd -District 2	Survey	0.4	No target weeds found	3
Vert Rd	Survey	0.3	No target weeds found	3
Wynn Ln	Survey	0.2	No target weeds found	3
Baskins Rd	Survey	0.4	No target weeds found	4
Elwha Rd	Survey	0.2	No target weeds found	4
Garling Rd	Survey	0.1	No target weeds found	4
Hudson Rd	Survey	0.4	No target weeds found	4
Hunt Rd	Survey	0.2	No target weeds found	4
Lookout Rd	Survey	0.4	N/A	4
Lower Dam Rd	Survey	0.5	N/A	4
Markuson Rd	Survey	0.2	No target weeds found	4
Nordstrom Rd	Survey	0.3	No target weeds found	4
North Bagley Creek Rd	Survey	0.1	No target weeds found	4
Old Rd	Survey	0.2	No target weeds found	4
Orvis St	Survey	0.1	No target weeds found	4
Prawn Rd	Survey	0.2	No target weeds found	4
Thors Rd	Survey	0.3	No target weeds found	4
Tongue Point Park Rd	Survey	0.6	N/A	4
Total: 162 roads		Total: 143.4 mi.		

¹Target Weed Species for Control – additional non-native, invasive species may be controlled as time and resources allow.

WEST CLALLAM COUNTY SELECTED ROADS

West Clallam County includes areas approximately west of Lake Crescent; the area includes Clallam Bay, Sekiu, Hoko-Ozette, Beaver, and Forks (reference Focus Area Maps 7-8). West Clallam County has a limited number of county roads and known infestations include knotweed, tansy ragwort, and Scotch broom. Roadside Scotch broom infestations impose significant negative impacts on adjacent forestry lands and have been elevated in priority at select locations in West Clallam County to reflect input and requests from local land managers.

Road Groups:

Clallam Co Restoration Project: roadsides adjacent to Clallam County environmental restoration or mitigation projects and may include native revegetation efforts; treatment is intended to contain infestations of non-native, invasive species to protect native plantings and function.

Source Cat. 1 Weeds: roadsides with the most extensive infestations of category 1 weeds; treatment is intended to contain and reduce infestations.

Isolated Cat. 1 Weeds: roadsides with isolated infestations of category 1 weeds either treated or surveyed in the past; treatment is intended to eradicate individual infestations or plants.

Forestry: roadsides that have known weed infestations and are used to access forestry managed timber lands; increasing potential for cross contamination

Pit Entrance: roadsides leading to County pits; treatment is intended to contain and protect county resources.

Public Request: roadsides with infestations requested by individuals or public agencies; treatment is intended to meet requests as time and resources allow.

Survey: roadsides that should be surveyed for any new weeds every 2-4 years; this group also contains roads that have not been surveyed for noxious weeds

Table 6. West Clallam County roads selected for herbicide treatment in 2025

Target Weed Species For Control – additional non-native, invasive species may be controlled as time and resources allow. Species listed in italics are unconfirmed infestations.

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Ballard Rd	Pit Entrance	0.3	yellow archangel, Scotch broom	1
Bear Creek Rd	Forestry	2.1	tansy ragwort	1
Bogachiel Way	Forestry	1.1	tansy ragwort	1
Coho Dr	Isolated Cat. 1 Weeds	0.3	tansy ragwort	1
Cooper Ranch Rd	Forestry	4.2	tansy ragwort, wild basil savory, herb Robert, Scotch broom	1
Dolly Varden Dr	Isolated Cat. 1 Weeds	0.2	tansy ragwort	1
Hermison Rd	Isolated Cat. 1 Weeds	0.6	Bohemian knotweed	1
Hoko-Ozette Rd	Forestry	21.4	Bohemian knotweed, herb Robert, Scotch broom, tansy ragwort, orange hawkweed	1

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Mary Clark Rd	Forestry	7.7	tansy ragwort, herb Robert, Scotch broom	1
Mina Smith Rd	Forestry	3.3	tansy ragwort, yellow archangel	1
Pillar Point Rd	Survey	0.3	No target weeds found	1
Quillayute Airport Rd	Isolated Cat. 1 Weeds	0.4	tansy ragwort	1
Quillayute Rd	Forestry	6.9	Scotch broom, tansy ragwort, common tansy	1
Rainbow Ave	Isolated Cat. 1 Weeds	0.4	tansy ragwort	1
Richwine Rd	Isolated Cat. 1 Weeds	0.6	tansy ragwort	1
River Park Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	1
Sekiu River Rd	Forestry	0.7	Bohemian knotweed, yellow archangel	1
Swan Bay Rd	Forestry	0.8	tansy ragwort	1
Undi Rd	Forestry	2.0	Bohemian knotweed, Himalayan blackberry	1
Wentworth Rd	Isolated Cat. 1 Weeds	1.2	tansy ragwort	1
Whitcomb-Diimmel Rd	Pit Entrance	1.5	Scotch broom, tansy ragwort	1
Wilson Rd	Isolated Cat. 1 Weeds	0.8	tansy ragwort	1
6th St	Survey	0.2	No target weeds found	2
7th St	Survey	0.3	No target weeds found	2
8th St	Survey	0.2	No target weeds found	2
Bloedel Blvd	Survey	0.2	No target weeds found	2
Bogachiel St	Isolated Cat. 1 Weeds	0.3	Bohemian knotweed, tansy ragwort	2
Calawah Way	Isolated Cat. 1 Weeds	1.4	tansy ragwort	2
Charley Creek Rd	Isolated Cat. 1 Weeds	0.7	yellow archangel	2
Commercial St	Isolated Cat. 1 Weeds	0.1	tansy ragwort	2
Division St	Isolated Cat. 1 Weeds	0.1	tansy ragwort	2
Duncan Rd	Survey	0.3	No target weeds found	2
Eagle Point Rd	Isolated Cat. 1 Weeds	1.3	Bohemian knotweed	2
East Lake Pleasant Rd	Isolated Cat. 1 Weeds	1.2	tansy ragwort	2
Elk Creek Ridge Rd	Forestry	0.2	tansy ragwort	2
Elk Valley Rd	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Front St	Isolated Cat. 1 Weeds	0.6	Bohemian knotweed, tansy ragwort	2
Funk & Ballards Aly	Survey	0.3	No target weeds found	2
Grant Rd – District 3	Survey	0.5	No target weeds found	2
Hilstrom Rd	Isolated Cat. 1 Weeds	0.6	Bohemian knotweed	2
Iverson Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	2
Kilmer Rd	Survey	0.8	No target weeds found	2
Lake Creek Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	2

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Lake Pleasant Park Rd	Survey	0.2	No target weeds found	2
Lucken Rd	Isolated Cat. 1 Weeds	0.2	tansy ragwort	2
Maxfield Rd	Forestry	0.8	tansy ragwort	2
Merchants Rd – Thimbleberry Ln to end	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Moriarty Rd	Forestry	1.1	tansy ragwort	2
Nichols Rd	Survey	0.6	No target weeds found	2
Old La Push Rd	Isolated Cat. 1 Weeds	0.4	Bohemian knotweed	2
Pavel Rd	Isolated Cat. 1 Weeds	1.3	tansy ragwort, Bohemian knotweed	2
Pit Ln	Pit Entrance	0.9	tansy ragwort	2
Rice St	Isolated Cat. 1 Weeds	0.1	tansy ragwort, orange hawkweed	2
Salmon Dr	Isolated Cat. 1 Weeds	0.3	tansy ragwort	2
Sekiu Airport Rd	Isolated Cat. 1 Weeds	0.3	tansy ragwort	2
Sekiu Heights Rd	Survey	0.2	No target weeds found	2
Shuwah Rd	Isolated Cat. 1 Weeds	0.5	Bohemian knotweed, yellow archangel	2
Sol Duc St	Survey	0.1	No target weeds found	2
Steelhead Ave	Isolated Cat. 1 Weeds	0.7	tansy ragwort, orange hawkweed	2
Taylor St	Isolated Cat. 1 Weeds	0.1	tansy ragwort, orange hawkweed	2
Trout Ave	Isolated Cat. 1 Weeds	0.5	tansy ragwort	2
Tyee Ridge Rd	Isolated Cat. 1 Weeds	0.3	orange hawkweed	2
Washington St	Isolated Cat. 1 Weeds	0.4	Bohemian knotweed, tansy ragwort	2
West Kellogg St	Survey	0.2	No target weeds found	2
West Lake Pleasant Rd	Isolated Cat. 1 Weeds	1.4	Bohemian knotweed	2
Wisen Creek Rd	Survey	0.4	No target weeds found	2
Conley Rd	Survey	0.5	No target weeds found	3
East Division St (Minnie Peterson Rd to 1601 E Division St)	Survey	0.8	No target weeds found	3
Elk Loop Dr	Isolated Cat. 1 Weeds	0.2	tansy ragwort	3
Frontier St	Survey	0.7	No target weeds found	3
Fuhrman Rd	Survey	0.3	No target weeds found	3
Gaydeski Rd	Isolated Cat. 1 Weeds	0.4	tansy ragwort	3
Green Rd	Survey	0.5	No target weeds found	3
Hoko Ln	Survey	0.1	No target weeds found	3
Kalawah St	Survey	0.1	No target weeds found	3
Kallman Rd	Survey	0.6	No target weeds found	3
Palmer Rd	Isolated Cat. 1 Weeds	0.5	tansy ragwort	3
Smith Rd	Survey	0.5	No target weeds found	3

Road Name	Group	Est. Road Miles	Target Weed Species For Control ¹	Treatment Priority
Vista Dr	Survey	0.3	No target weeds found	3
Vista Ln	Survey	0.1	No target weeds found	3
Walgren Rd	Survey	0.3	No target weeds found	3
Walter Way	Survey	0.2	No target weeds found	3
Weel Rd	Survey	0.3	No target weeds found	3
Erickson Rd	Survey	0.1	No target weeds found	4
Leyendecker Rd	Survey	0.1	No target weeds found	4
Total: 85 roads		Total: 85.6 mi.		

¹Target Weed Species for Control – additional non-native, invasive species may be controlled as time and resources allow.

COUNTY PITS

An important component of the IWM Program is prevention, such as maintaining weed free rock sources. County-owned rock sources and storage pits sites are routinely surveyed for weeds and specific management strategies are tailored to match each pit's county use and needs. Most are gated, with no public access.

The 2025 Work Plan includes all County-owned pits as potential locations for weed control. The County Pits vary in size, activity, and weed pressures. Table 7 lists the county pits in alphabetical order and includes the target weed species for control and approximate pit location. Control methods may include physical or chemical methods, but more likely a combination of both, depending on weed species, area, and available resources. Cultural practices, such as reseeding with locally sourced plant material will be pursued as materials become available and suitable pit locations identified.

Table 7. County Pit Sites for treatment in 2025

Pit Name	Location	Target Weed Species For Control ¹
Blue Mountain Transfer Station: 6.8 ac	Blue Mountain Rd, Port Angeles	bull thistle, Canada thistle, meadow knapweed, poison hemlock, reed canary grass, Scotch broom
Blyn Pit: 18.65 ac	Woods Rd, Blyn	Bohemian knotweed, butterfly bush, common teasel, everlasting pea vine, field bindweed, hairy nightshade, herb Robert, Himalayan blackberry, poison hemlock, Scotch broom, tansy ragwort
Clallam Bay Storage Yard: 4.65 ac	Frontier St, Clallam Bay	Himalayan blackberry, reed canary grass
District 1 Shop: 4.15 ac	W Washington St, Sequim	bull thistle, Canada thistle, Himalayan blackberry, meadow knapweed, spotted knapweed, St. John's wort
District 2 Shop: 5.22 ac	W Lauridsen Blvd, Port Angeles	herb Robert, meadow knapweed, poison hemlock, Scotch broom
Forks Pit: 21.36 ac	Pit Ln, Forks	Bohemian knotweed, tansy ragwort, bull thistle, Canada thistle, Himalayan blackberry
Herrick Pit: 11.5 ac	Herrick Rd, Port Angeles	bull thistle, Canada thistle, everlasting pea vine, herb Robert, Himalayan blackberry, meadow knapweed, Scotch broom
Hogback Pit: 1.76 ac	Cays Rd, Sequim	common tansy, tree lupine
Hoko-Ozette Rd MP 4.5: 1.0 ac	MP 4.5 Hoko-Ozette Rd, Clallam Bay	herb Robert, orange hawkweed, tansy ragwort, Scotch broom
Hoko-Ozette Rd MP 10: 2.85 ac	MP 10 Hoko-Ozette Rd, Clallam Bay	herb Robert, tansy ragwort
Hoko-Ozette Rd MP 13: 2.98 ac	MP 13, Hoko-Ozette Rd, Clallam Bay	herb Robert, tansy ragwort
Hwy 101 Storage Yard: 4.03 ac	Hwy 101/Old Olympic Hwy, Port Angeles	herb Robert, poison hemlock, Himalayan blackberry
Kirner Pit: 37.37 ac	Kirner Rd, Sequim	bull thistle, common chicory, common periwinkle, common teasel, Himalayan blackberry, lesser burdock, meadow knapweed spotted knapweed, poison hemlock, Scotch broom, tansy ragwort, tree lupine, yellow archangel, moleplant

Pit Name	Location	Target Weed Species For Control ¹
La Push "Ballard" Pit: 8.11 ac	Ballard Rd, Forks	Canada thistle, cut leaf blackberry, everlasting pea vine, Himalayan blackberry, Bohemian knotweed, Scotch broom, tansy ragwort, yellow archangel
Lake Creek Pit: 11.47 ac	Highway 101 MP 198.7, Beaver	bull thistle, Canada thistle, cut leaf blackberry, Himalayan blackberry, Bohemian knotweed, Scotch broom, tansy ragwort
Little River Pit: 5.05 ac	863 Little River Rd, Port Angeles	bull thistle, Canada thistle, herb Robert, meadow knapweed, Scotch broom, everlasting pea vine, tansy ragwort
Lower Elwha Pit: 6.22 ac	2045 Lower Elwha Rd, Port Angeles	bull thistle, Canada thistle, poison hemlock, reed canary grass, Scotch broom
McInnes Pit: 5.5 ac	Vistas Drive, Sequim	Bohemian knotweed, wild mustard, bull thistle, burr chervil, Canada thistle, common teasel, herb Robert, field bindweed, Himalayan blackberry, Italian thistle, meadow knapweed, poison hemlock, reed canarygrass, Scotch broom, spotted knapweed, common tansy
Morse Creek Pit: 33.68 ac	Gravel Pit Rd, Port Angeles	Bohemian knotweed, bull thistle, Canada thistle, common tansy, common teasel, herb Robert, Himalayan blackberry, poison hemlock, reed canary grass, Scotch broom
Piedmont Pit: 21.22 ac	Joyce-Piedmont Rd, Joyce	bull thistle, Canada thistle, common teasel, herb Robert, meadow knapweed, Scotch broom, tansy ragwort
Place Pit: 4.94 ac access shared with WSDOT	Place Rd, Port Angeles	Bohemian knotweed, bull thistle, butterfly bush, Canada thistle, common teasel, cut leaf blackberry, everlasting pea vine, herb Robert, Himalayan blackberry, meadow knapweed, oxeye daisy, reed canary grass, Scotch broom, tree lupine
Quillayute Pit: 20.75 ac	Quillayute Rd, Forks, WA	Bohemian knotweed, Himalayan blackberry, Scotch broom, tansy ragwort
Ranger Pit: 19.3 ac	Place Rd, Port Angeles	Bigleaf periwinkle, Bohemian knotweed, bull thistle, butterfly bush, Canada thistle, common teasel, cut leaf blackberry, everlasting pea vine, field bindweed, herb Robert, Himalayan blackberry, meadow knapweed, reed canary grass, Scotch broom, tansy ragwort, tree lupine
Sequim Storage Yard: 2.57 ac	W Washington St, Sequim	wild mustard, common chicory, common teasel, cut leaf blackberry, Himalayan blackberry, meadow knapweed, oxeye daisy, poison hemlock, reed canary grass, spotted knapweed, spurge laurel
Umbrella Creek Pit: 14.62 ac	Hoko-Ozette Rd, Clallam Bay	Bohemian knotweed, cut leaf blackberry, herb Robert, Himalayan blackberry, Scotch broom, tansy ragwort
Whitcomb-Diimmel Pit; 15.74 ac	Whitcomb-Diimmel Rd, Forks	Bohemian knotweed, bull thistle, Canada thistle, common teasel, Himalayan blackberry, Scotch broom, tansy ragwort, St. John's wort

¹Target Weed Species for Control – additional non-native, invasive species may be controlled as time and resources allow.

COUNTY SPECIAL SITES

The Road Department has land management responsibilities for several non-roadside properties called “Special Sites”. Locations listed in Table 8 are listed in alphabetical order and include the location of the site, a description and target weed species for control. The list includes a variety of sites, including designed features with specific functions and/or management requirements. The sites have been prioritized for weed management and control may include the use of herbicides. Other Road Department-managed lands that are encountered during other weed control projects that are not listed in Table 8 may be included as EDDR site.

Table 8. County Special Sites selected for treatment in 2025

Site Name	Location	Description	Target Weed Species For Control ¹
Black Diamond Native Planting	Adjacent to 223 Black Diamond Rd, Port Angeles	Native planting maintenance	Canadian thistle, Scotch broom
Blake Sand & Gravel Ridge	Cays Rd, Sequim	Ridge and slope northwest of Hogback Pit	Italian thistle, poison hemlock
Carlsborg Road Fire District	Carlsborg Rd, Carlsborg	Large county parcel adjacent to fire maintenance facility, school, and housing development.	poison hemlock
Cays & Lamar Intersection	Cays Rd & Lamar Rd, Sequim	County-owned parcel between Cays Rd and Lamar Rd	Italian thistle
Deer Park Overpass and Rest Area	Deer Park Loop, Port Angeles	Construction and slope stabilization project; control of non-native, invasive weeds to protect constructed slopes and adjacent areas to prepare for revegetation. Control of <u>vegetative</u> invasive blackberries ONLY.	common teasel, cut leaf blackberry, Himalayan blackberry, meadow knapweed, Scotch broom
Northern levee trail Parking Lot	Towne Rd, Sequim	School House parking lot with access to multi-use trail along Dungeness Floodplain restoration project	poison hemlock
Dungeness River Bridge	Old Olympic Highway, Sequim	County-owned parcel with small foot trail down to the Dungeness River under the bridge	Bohemian knotweed, butterfly bush, Canada thistle, Himalayan blackberry, Scotch broom
Dungeness Trails	River Rd, Sequim	County-owned site currently managed by Clallam County Public Works; potential future site of Dungeness Reservoir	Meadow knapweed, poison hemlock, common teasel
East Ennis Creek	E Ennis Creek Rd, Port Angeles	County owned property on E Ennis Creek Rd	Yellow archangel

Site Name	Location	Description	Target Weed Species For Control ¹
East Greentree Ln	E Greentree Ln, Port Angeles	County owned property on E Greentree Ln	Yellow archangel
Edwards Rd - Clallam County Storage Yard	Edwards Rd, Port Angeles	Clallam County Search and Rescue, Radio Maintenance Storage yard	Meadow knapweed
Gasman Detention Pond	Gasman Rd, Port Angeles	Stormwater detention site	Scotch broom
West Elwha Trailhead	End of Herrick Rd, Port Angeles	Reroute of West Elwha trailhead	spurge laurel
Hoko-Ozette Culvert Installation MP 8.8	Hoko-Ozette Rd, Sekiu	Culvert installation and revegetation project; control of non-native, invasive weeds and support the establishment of native plantings at constructed culvert locations.	Scotch broom
Jamestown Beach Bank Stabilization	Jamestown Rd, Sequim	Revegetation and bank stabilization project; control of non-native, invasive weeds to support the establishment of native plantings.	Survey data not available
Kugel Creek Culvert Installation	Cooper Ranch Rd, Sappho	Culvert installation and revegetation project; control of non-native, invasive weeds and support the establishment of native plantings at constructed culvert locations.	herb Robert
Lower Dam Interpretive Center	Lower Dam Rd, Port Angeles	Trail head, parking lot, and interpretive center;	bull thistle, mint
Lower Elwha Rd Culvert Installation	Lower Elwha Rd, Port Angeles	Culvert installation and revegetation project; control of non-native, invasive weeds and support the establishment of native plantings at constructed culvert locations.	Canada thistle, evergreen blackberry, Himalayan blackberry, Scotch broom
Marsden Rd Culvert Installation	Marsden Rd, Port Angeles	Culvert installation and revegetation project; control of non-native, invasive weeds and support the establishment of native plantings at constructed culvert locations.	Survey data not available
McDonald Creek Bridge	4100 - 4132 Old Olympic Highway, Agnew	Bridge construction and revegetation project; control of non-native, invasive weeds and support the establishment of native plantings at constructed slopes.	Canada thistle, cut leaf blackberry, everlasting pea vine, herb Robert, Himalayan blackberry
Mt. Pleasant Storm water Raingarden	Mount Pleasant Rd, Port Angeles	Constructed storm water raingarden; control of non-native, invasive weeds to support the establishment of native plantings.	Survey data not available

Site Name	Location	Description	Target Weed Species For Control ¹
O'Brien ROW parcel	O'Brien Rd, Port Angeles	Large ROW parcel, adjacent to Erving Jacobs intersection. Public request.	Canada thistle
Olympic Discovery Trail	County wide	Multi-user transportation/recreational trail; control of Category 1 species and Category 2 species as time and resources allow; control of <u>vegetative</u> invasive blackberries ONLY.	Canada thistle, common teasel, European hawkweed, herb Robert, meadow knapweed, poison hemlock, spotted knapweed, Scotch broom
	Berm on Old Olympic Hwy east of Lewis Rd, Agnew	Multi-user transportation/recreational trail; revegetation project; control of non-native, invasive weeds and support the establishment of native plantings at constructed slopes.	wild mustard, bull thistle, burr chervil, Canada thistle, reed canary grass
	Dean Creek bridge	Multi-user transportation/recreational trail; from 1.0 mi west of bridge to bridge	spurge laurel, Bohemian knotweed
	Gossett Rd to Gossett Bridge, Joyce	Multi-user transportation/recreational trail; revegetation project; control of non-native, invasive weeds and support the establishment of native plantings at constructed slopes.	Canada thistle, herb Robert, meadow knapweed, reed canary grass, tansy ragwort
	Hwy 101 crossing near Sol Duc	Multi-user transportation/recreational trail	European hawkweed
	Priest Rd, Sequim	Multi-user transportation/recreational trail; access to Dungeness River Nature Center	poison hemlock
	West Sequim Bay Rd to Whitefeather Way	Multi-user transportation/recreational trail; in conjunction with Spoke Folk bicycle club	Himalayan blackberry
Olympic Wetland Mitigation Site	91 E Hwy 101, Port Angeles	Wetland mitigation project: control of non-native invasive weeds and support the establishment of native plantings.	Canada thistle, Scotch broom, sulfur cinquefoil, reed canary grass
Pearce Rd Culvert Installation	Pearce Rd, Port Angeles	Culvert installation and revegetation project: control of non-native, invasive weeds to support the establishment of native plantings at constructed culvert locations.	Canada thistle, mint, Scotch broom
Pillar Point Bank Stabilization	Pillar Point Recreation Area, Pysht	Revegetation and bank stabilization project; control of non-native, invasive weeds to support native plantings.	perennial sow thistle, Scotch broom, tansy ragwort
Towne Rd berm (southern end of levee)	Towne Rd, Sequim	New berm where levee meets road; MANUAL REMOVAL ONLY	Common teasel
Towne Rd levee pedestrian path	Towne Rd, Sequim	Pedestrian path along Towne Rd levee, including levee slopes	Poison hemlock, wild mustard, everlasting peavine, Canada thistle, other non-native weeds that are spreading from levee sides into path

Site Name	Location	Description	Target Weed Species For Control ¹
Ward Bridge Restoration	Woodcock Rd, Sequim	Bridge construction and revegetation project; control of non-native, invasive weeds and support native plantings.	herb Robert, poison hemlock
Weel Road Bank Stabilization	Weel Road, Clallam Bay	Revegetation and bank stabilization project; control of non-native, invasive weeds to support native plantings.	Survey data not available
Voice of America Parking lot	Dungeness Recreation Area	Large pullout at mile 0.25	Poison hemlock

¹Target Weed Species for Control – additional non-native, invasive species may be controlled as time and resources allow.

FOCUS AREA MAPS OF SELECTED ROADS

The following maps display the County roads, pits, and Special Sites selected for control in 2025. All selected roads are color coded to the full extent of County management, but treatments may be limited to weed locations and, in many cases, only at isolated patches of Category 1 weeds. Selected roads are highlighted by color to represent their treatment priority.

Priority 1 selected roads are color coded in **green** and are intended to be completely treated for target species

Priority 2 selected roads are color coded in **yellow** and are intended to be treated for target species in conjunction with adjacent roads or activities

Priority 3 selected roads are color coded in **orange** and are intended to be treated or surveyed only as time and resources allow

Priority 4 selected roads are color coded in **blue** and are intended to be put onto a 4-year maintenance regime where no treatment is necessary

County-owned Special Sites and pits are included in the focus area maps. 2024 roadside treatment points, when taken, have been included for reference.

Overview of Clallam County

Map 1. Clallam County Roads District Overview

East Clallam County

Map 2. Miller Peninsula – Blyn Treatment Area

Map 3. Sequim – Dungeness Valley Treatment Area

Map 4. South Sequim – Carlsborg Treatment Area

Central Clallam County

Map 5. Agnew – East Port Angeles Treatment Area

Map 6. Port Angeles – Elwha Treatment Area

Map 7. Joyce – Lake Sutherland Treatment Area


West Clallam County

Map 8. Sekiu – Clallam Bay – Ozette Treatment Area

Map 9. Sappho – Sol Duc Valley Treatment Area

Map 10. Forks – Quillayute Rd Treatment Area

Clallam County Roads District Overview




Miller Peninsula-Blyn Treatment Area




Sequim - Dungeness Valley Treatment Area



South Sequim - Carlsborg Treatment Area



Agnew-East Port Angeles Treatment Area




Port Angeles - Elwha Treatment Area




Joyce - Lake Sutherland Treatment Area



Sekiu-Clallam Bay-Ozette Treatment Area



Sappho-Sol Duc Valley Treatment Area



Forks-Quillayute Rd Treatment Area



HERBICIDE PRODUCT LIST

Products chosen for this program are effective on known roadside and pit weeds, offer the greatest weed selectivity, maximize worker and public safety (no wait, access when the spray has dried), and pose the lowest risk for wildlife and the environment. See Appendix B for greater detail of risk analysis, product selection process and description of application methods. Because the preferred provider may vary from year to year; different brand names listed in the following tables with the same active ingredient may be substituted. New products or different formulations with the same active ingredient that are more user-friendly or environmentally friendly and cost-beneficial will be substituted as they become available.

Clallam County identifies the following products for targeted herbicide applications:

- AquaNeat® (aquatic formulation glyphosate)
- Element 3A® (aquatic formulation triclopyr TEA)
- Fusilade II® (fluazifop-P)
- Garlon 3A® (aquatic formulation triclopyr TEA)
- Garlon 4® (triclopyr BEE)
- HighNoon® (aquatic formulation aminopyralid and florypyrauxifen)
- Milestone® (aquatic formulation aminopyralid)
- Oust XP® (sulfometuron-methyl)
- Plateau® (imazapic)
- Polaris® (aquatic formulation imazapyr)
- Ranger Pro® (glyphosate)
- Transline® (clopyralid)
- Vastlan® (aquatic formulation triclopyr choline salt)
- WeeDestroy AM-40® (aquatic formulation 2,4-D)

APPENDICES

Appendix A: Ordinance 923 – Integrated Weed Management

Appendix B: Non-target Impacts and Risk Assessment, Herbicide Selection Process, Product List, and Application Methods

Appendix C: Sample Record Keeping Forms

Appendix D: Sample Press Release and Public Notice

Appendix E: Sample Herbicide Notice

Appendix F: Owner Will Control Packet

Appendix G: Adopt-a-Patch Permit

Appendix H: Historical Data for Known Roadside Weed Locations

Appendix I: References

Ordinance 923

AN ORDINANCE creating a chapter titled "*Integrated Weed Management*"

BE IT ORDAINED BY THE BOARD OF CLALLAM COUNTY COMMISSIONERS:

Section 1. Section .010, Purpose, is created to read as follows:

The purpose of this chapter is to establish an integrated weed management approach that aids Clallam County in keeping county owned or managed lands, including its improved and unimproved right-of-ways, quarries, parks and other county lands safe and operating at an efficient level of service while meeting its legal obligations and stewardship responsibilities to control noxious weeds.

Furthermore, it is the legislative body's intent that Clallam County's departments, employees, and designees have the ability and flexibility to address and combat the spread of noxious weeds in our community.

Integrated weed management should promote desirable vegetation and remove noxious and invasive plant species of special concern, through comprehensive, strategic, environmentally responsible, and cost effective methods, including mechanical, biological, cultural, chemical, and prevention control methods as needed.

This chapter is subject to applicable federal and state laws as adopted and hereafter amended; and supersedes the Clallam County Roadside Vegetation Management Policy and any subsequent resolutions relating thereto.

Section 2. Section .020, Applicability, is created to read as follow:

The provision of this chapter applies to any Clallam County department, employee, or designee that may perform work within county right-of-ways, parks or other county owned or managed lands with regard to weed control as defined in this chapter.

Inclusion of this chapter in this specific title is not intended to limit its scope and application to all county owned or managed lands.

Section 3. Section .030, Definitions is created to read as follows:

For purposes of this section, the following definitions apply:

(1) "Biological control" means using living organisms that suppress the host plant. Insects, diseases, and foraging animals are examples of biological controls.

(2) "Chemical control" means using conventional or natural herbicides to eliminate noxious weeds or additional invasive, non-native plant species that present a special concern within a specified area. It does not mean treatments for general vegetation management.

(3) "Cultural control" means enhancing the vigor of desirable plants which may crowd-out or prevent weed infestations. Hydroseeding or planting low-growing, self-maintaining shrubs are examples of cultural practices.

(4) "Feasible" means a control method or combination thereof that is capable of being carried out or achieving a goal. This includes consideration of such factors as plant biology, site characteristics, scope of the problem, and available resources.

(5) "Integrated Weed Management (IWM)" means a coordinated decision making process that uses the most appropriate weed management methods and strategies, along with a monitoring and evaluation system, to achieve weed management goals and objectives in an environmentally and

economically sound manner. It allows for the use of mechanical, biological, cultural, chemical, and prevention control methods.

(6) "Invasive plant" means an introduced, non-native plant that is aggressive and causes economic loss and adverse effects to agricultural, natural and human resources.

(7) Invasive plant of special concern means an invasive plant identified as a threat by the US Department of Agriculture, the Washington Departments of Agriculture and Ecology, or the Washington Invasive Species Council.

(8) "Mechanical, manual control" means cutting or cultivating in a manner to reduce or slow undesirable plant growth (*i.e.*, using hand labor, mowers, graders, and ditching equipment).

(9) "Planning and prevention" means creating design standards that enhance or complement other control methods (such as slope grade within mower reach) and policies or standard operating procedures such as equipment cleaning or weed free material standards that prevent contamination or spread of noxious or non-native invasive species.

(10) "Weeds" means both noxious weeds which are so designated under State law, and additional non-native invasive plants of special concern that have been added to a County specific list of plants targeted for control and elimination.

(11) "Work Plan" means a plan prepared annually by the Clallam County Road Department or any other county department with land management responsibilities, and approved by Clallam County Noxious Weed Control Board prior to the commencement of seasonal weed management.

Section 4. Section .040, Integrated Weed Management Plan, is created to read as follows:

Each department with land management responsibilities, such as the Road Department, shall develop and implement an Integrated Weed Management Plan (hereinafter "IWM Plan" or "work plan"), that is specific to land under their management. Each work plan will be reviewed annually, and amended as needed, in concert with and approved by the Noxious Weed Control Board.

Each IWM Plan shall contain a list of priority species and must include the Clallam County Noxious Weed List. Other invasive species may be added as resources allow and according to the level of threat posed. Work plans shall also contain provisions for early detection/rapid response (EDRR) that allows additional weeds, locations, and techniques to be added to a work plan as necessary should new infestations of high priority weeds be discovered during the course of the treatment season.

IWM Plans shall contain detailed information about departmental practices and standards, including but not limited to:

(1) Location of high-priority weed infestations, sensitive areas, and other areas with special management considerations.

(2) Guidelines and prescriptions for best management practices in dealing with weed problems and opportunities, including planning and prevention measures.

(3) Monitoring protocols used to evaluate infestation status and treatment effectiveness, the results of which are publically available and used to inform subsequent work plans.

(4) Creation of an outreach and education component including coordinated volunteer opportunities as time and resources allow.

A copy of each IWM Plan will be maintained by the Department for which it is developed and the Noxious Weed Control Board.

Section 5. Section .050, Citizen Option, is created to read as follows:

(1) Property owners that do not wish herbicides to be applied to eliminate noxious weeds or other invasive weeds of special concern to road right-of-ways or county managed lands directly bordering their property may enter into an annual "owner will control" agreement with Clallam County. Those with current agreements are allowed to post the county adjacent boundary as no spray areas.

(2) Under an "owner will control" agreement, the property owner must undertake specific measures, prescribed within the agreement that will ensure the timely and effective control and reduction of target weed species within the right-of-way or within a proscribed area specifically identified in the agreement


(3) If the property owner fails to effectively control or reduce targeted weed species as agreed, then Clallam County will issue a single written warning. If the problem persists after 10 days from when the written warning was sent, then Clallam County reserves the right to void the agreement upon written notice to the property owner.

(4) If any agreement is subsequently voided, Clallam County may immediately proceed with any planned weed control deemed necessary.


(5) A property owner may re-apply for a new agreement in a subsequent season; however, the County reserves the right to deny said application if there is a documented history of failure to adhere to the terms of such agreements or history of voided agreements.

ADOPTED this tenth day of January 2017


BOARD OF CLALLAM COUNTY COMMISSIONERS



Mark Ozias, Chair



Randy Johnson



Bill Peach

ATTEST:



Loni Gores, Clerk of the Board

APPENDIX B: NON-TARGET IMPACTS AND RISK ASSESSMENT, HERBICIDE SELECTION PROCESS, PRODUCT LIST, AND APPLICATION METHODS

Non-Target Impacts and Risk Assessment

Not only must a weed control strategy be effective and efficient, but it must consider potential adverse impacts to non-target plant and animal species and include measures to mitigate those impacts to the greatest extent possible.

Any potential impacts to humans, pets, livestock, wildlife, desirable plants and the environment from noxious weed and invasive plant removal are of concern to the project managers. Every control method has benefits and costs. For example, the disturbance caused by workers and mowing or excavation equipment which allows weeds to proliferate can be more significant than impacts from herbicides. Hand removal may result in trampling and soil disturbance. Although all control methods pose some level of risk, potential risks associated with herbicide use will receive the greatest scrutiny. Best management practices that reduce or mitigate potential herbicide impacts to non-target organisms will be incorporated into all aspects of the work plan.

An assessment of risk involves understanding the toxicity and likely exposure paths for various organisms that may be exposed to an herbicide. Risk assessments are used by project managers to identify those exposures that might be problematic. The project manager then uses this information to decide whether herbicides can be used without undue risk and to develop mitigation actions to reduce risks. A critical component of properly applying this risk assessment process is creating the institutional conditions for obtaining and retaining project managers who either receive or have had significant training and depth of experience to make and apply these decisions.

Several concepts are important in minimizing adverse effects. At a minimum, herbicide users should be familiar with:

1. The relative risk posed by the herbicide to the applicator and general public, and the anticipated exposure scenarios.
2. The types of wildlife and vegetation present, including endangered species. The invasive weed manager should learn enough about each species (life cycle, breeding habitat, food supply, shelter needs, etc.) to avoid impacts.
3. The relative risk posed by the herbicide to different wildlife and plant taxa that may be present and the anticipated exposure scenarios. Consideration should be given not only to the active ingredient, but also other compounds added to an herbicide formulation or added to the “tank mix” to be applied, such as surfactants.
4. The relative persistence of the herbicide in the environment, primarily in soil. Herbicide persistence is measured in terms of “half-life.” One half-life is the amount of time it takes for the herbicide to break down to 50% of its original concentration in soil or water. Generally, it takes five half-lives for more than 97% of the herbicide to be fully degraded. Herbicide persistence is discussed in more detail in Appendix B.
5. The mobility of the herbicide in runoff water. Off-site movement in surface water and leaching to groundwater are both primarily influenced by the amount applied, the herbicide’s water solubility and its tendency to adsorb to soils. Factors affecting herbicide mobility are discussed in more detail in Appendix B.

Important background information regarding the types of animals that may be impacted by noxious weed and invasive plant control in a roadside setting has been synthesized from the California Invasives Species Council (Cal-IPC) 2015 manual titled, *Best Management Practices for Wildland Stewardship: Protecting Wildlife When Using Herbicides for Invasive Plant Management* and presented below. The full document is available at www.cal-ipc.org. Such information is vital to making informed decisions on ways to mitigate or avert potential

effects where possible; especially when making control choices about herbicide selection, application methods and timing. Although the Cal-IPC manual's focus is on wildlands, many of the same types of animals may be found living near, if not on, county roadsides and should therefore be considered.

Organisms that are endangered or under threat of becoming endangered receive special protection under The Endangered Species Act. An Information for Planning and Conservation (IPaC) Trust Resource report of threatened and endangered plant and animal species found in Clallam County was provided by the US Fish and Wildlife Service. These species are included in Table 10.

Insects

Insects are a diverse class of animals that are part of the food web on which many vertebrate species depend. Butterflies, bees, wasps and even mosquitoes pollinate plants that then provide fruits and seeds for other animals. Flies and beetles eat rotting debris, which helps recycle nutrients in the ecosystem. Aphids and many other soft-bodied insects suck the juices of plants and are themselves a high-protein food for other insects, reptiles, amphibians, birds, and mammals.

Most insects are so small and so intimately connected to vegetation that it is difficult to avoid spraying them directly, along with the invasive plants being treated. Honeybees are routinely tested for sensitivity to herbicides and are broadly representative of other insects. While most herbicide active ingredients used in wildland weed management pose very low toxicological risks to invertebrate species, some of the inert ingredients in formulated herbicide products may pose a greater risk. For example, some oil-based emulsifiable concentrate formulations may be harmful to soft-bodied adults or larval insects like aphids or caterpillars. The Taylor's checkerspot butterfly is the only endangered insect listed in Clallam County. No roadside habitat for the Taylor's checkerspot butterfly has been identified.

Reptiles and Amphibians

Lizards, snakes, turtles, frogs, newts, and salamanders are frequently residents of areas where invasive plant management is planned. These species can be exposed to herbicides through direct sprays and spray drift, and through consuming herbicide-contaminated water, prey, or plants. Amphibians may be especially vulnerable, since they spend a portion of their life cycle as aquatic organisms and often only need small puddles or seasonal streams for growth. The inert ingredients in a formulated herbicide product may be as important to evaluate as the active ingredient in terms of the risk they pose to amphibians. No endangered or threatened reptiles or amphibians have been identified in Clallam County.

Fish and Aquatic Invertebrates

Fish and aquatic invertebrates are often more sensitive to herbicides than terrestrial animals because of their physiology or the increased exposure potential that may result from herbicide movement into aquatic sites. Aquatic species can be exposed to herbicides through direct spray, spray drift, spills or surface runoff. Though few commonly used herbicide active ingredients are highly acutely toxic to aquatic organisms, toxic effects can result from the exposure to other ingredients in formulated products, such as surfactants. With the current suite of herbicides typically used in invasive plant management, bioaccumulation of herbicides in fish tissue is not a problem, since these herbicides are typically metabolized and/or excreted fairly quickly. A number of fish species (salmonids and trout) found in Clallam County are listed as threatened or endangered and many creeks and rivers are the subject of habitat restoration projects intended to help restore these stocks to healthy population levels.

Mammals

Deer, coyotes, mountain lions, wood rats, gophers, and mice are just a few of the mammals that may populate or feed on animals that populate typical roadsides. Pets, such as dogs and cats, with their owners or

wandering freely, might be exposed in a more limited manner. Animals may be exposed to herbicides through contaminated food or water, as well as direct sprays, spray drift, and contact with treated vegetation. The toxicity of herbicides to mammals has been better studied than for most other species because they are used as surrogates for human toxicity assessments. Studies on mammals allow for evaluation of a wide variety of parameters, including reproductive, developmental, and neurological effects in exposed populations, as well as effects on blood chemistry, organ weights, and body weight gain or loss.

The most abundant mammals on a typical roadside area are rodents. They are small enough and abundant enough that they may be directly sprayed or exposed to drift during an herbicide application, particularly with ground spray equipment.

Deer and other herbivores may browse on treated vegetation. Once the vegetation is dead, it becomes less attractive to eat; however, in situations where a selective herbicide is used that kills only broadleaf plants or only grass plants, the treated, but unaffected plant species may pose a dietary exposure risk.

Fishers, while not currently listed as an endangered species, have received special management consideration and have been reintroduced into Clallam County in Olympic National Park. No county roadside habitat has been identified.

Birds

Potentially impacted birds include large carnivorous birds like hawks or ospreys, herbivorous species like geese and ducks, small insectivorous birds, and small fruit and seed-eating birds. All of these species can be exposed to herbicides through their food and drinking water. The highest risks are typically for birds eating sprayed vegetation since that is often the target of the application, and the likelihood of being exposed is higher than for those species eating contaminated prey. In general, the herbicides used to control invasive plants do not pose significant acute toxicity risks to birds when used under typical use scenarios; however, less is known about chronic and reproductive effects. To minimize risk, applications during nesting season should be avoided if possible. Several federally listed bird species are found in Clallam County, but there are no habitat listings for county right-of-way.

Plants

All types of plants may be affected by weed control activities. Because herbicides are designed to kill plants, an applicator's ability to distinguish desirable plants from weeds is critical. Certain native plant species are protected under state or federal laws. The most current data set (as a GIS shapefile) was obtained from the Natural Heritage Program which is managed by the Washington State Department of Natural Resources. It contained general locational information of rare, threatened, and endangered plant species. It was reviewed for species and sites that warrant special management consideration on Clallam County roadsides. Pink fawn lily, which is noted as sensitive (a non-regulatory status) was found in the vicinity of six county roadsides. No noxious weed infestations have yet been documented near these pink fawn lily sites, but all shall be noted and continue to be under special consideration. Whitebark pine which is a candidate for federal listing is known to exist in Clallam County, but no sites have been identified on county right of way. No rare, endangered or threatened species were identified on county roadsides in the DNR Natural Heritage Program dataset.

Table 9. Species in Clallam County with potential for special management consideration

COMMON NAME	POPULATIONS PRESENT /HABITAT	POPULATION IDENTIFIED ON COUNTY ROADSIDE	LISTING STATUS
PLANT			
Pink fawn lily	Yes	In vicinity of Walgren Rd, Grant Rd, Pavel Rd, River Breeze Wy, West Lake Pleasant Rd, Hoko-Ozette Rd	State-Sensitive, (non-regulatory)
Whitebark pine	Yes	none	Fed Candidate
BIRD			
Streaked Horned Lark	Yes	none	Fed-Threatened
Marbled murrelet	Yes/plus habitat overlap	none	Fed-Threatened
Northern spotted owl	Yes/plus habitat overlap	none	Fed-Threatened
Short-tailed Albatross	Yes	none	Fed-Endangered
Yellow-billed Cuckoo	Yes	none	Fed-Threatened
FISH			
Bull Trout	Yes/plus habitat overlap	Indirect	Fed-Threatened
Dolly Vardon	Yes	Indirect	Fed-Threatened
INSECT			
Taylor's checkerspot butterfly	Yes-FS, private, ONP, DNR/ plus habitat overlap	None - possible potential habitat?	Fed-Endangered
MAMMAL			
Fisher	Yes, reintroduced in ONP	None	Fed-Threatened
HABITAT OVERLAP ONLY			
Chinook	Habitat designation	Indirect	Threatened
Chum	Habitat designation	Indirect	
Sockeye	Habitat designation	Indirect	
Killer whale	Habitat designation	Indirect	

Risk Charts

The herbicide risk charts, tables, and text that follow have been reproduced with permission from the publication: Cal-IPC. 2015 *Best Management Practices for Wildland Stewardship: Protecting Wildlife When Using Herbicides for Invasive Plant Management*. Cal-IPC Publication 2015-1. California Invasive Plant Council, Berkeley, CA. The charts include the most common herbicides used by wildland managers for invasive plant management and include those chosen for use on Clallam County roadsides (see Table 13 in Appendix B). Fluazifop, which would be allowed for use under this plan, was not included in the Cal-IPC risk charts because the data needed to conduct the analysis was not available at the time the risk charts were completed.

Wildlife

The risk charts provide information on the comparative risk of each herbicide to each type of wildlife from selected exposure scenarios. A summary of the methods used to generate these charts follows and refers the reader to the primary sources for more detail. Each chart summarizes potential risk for a specific exposure scenario and is based on a risk

assessment model developed by the USFS. See the spreadsheet of calculations on the PRI website for detailed information on risk charts. Using the spreadsheet, you can modify application rates to assess changes in risk profiles. It is important to note that many of the scenarios are “worst case” and do not represent typical real-world situations. The assumptions for each scenario, with a description

about how they relate to typical real-world situations are listed on the risk charts.

Risks that fall outside an acceptable zone should prompt the land manager to consider steps to mitigate the risk.

Risk to Wildlife Depends on Both Toxicity and Exposure

Risks to wildlife are dependent on the herbicide's toxicity to that particular taxonomic group and the animal's exposure to the herbicide. Toxicity is described using Toxicity Reference Values (TRVs), which represent the dose of herbicide generally assumed to be without adverse effects. Lower TRVs indicate a more toxic herbicide for the particular taxonomic group. The TRVs used to develop the risk charts for the different wildlife taxa are summarized in below.

An important determinant of exposure is the herbicide application rate. For the risk charts, the application rates were set to half of the maximum application rate as indicated on the herbicide's product label. This "half- max" application rate was used to better approximate typical wildland herbicide applications. For example, invasive plant management typically involves portions of acres to be spot treated, but not entire acres. Alternatively, entire acres might be treated via broadcast spray, but at rates below maximum allowable rates. Since application rate is directly proportional to risk, the risk values at maximum application rates would simply be twice the values shown in the charts (likewise, lower rates would have proportionally less risk)—with the exception of spills, where application rate is not relevant. Table 11 provides the application rates used to estimate exposure for each herbicide in terms of pounds of the active ingredient (or the acid equivalent of the active ingredient) and the equivalent rate per acre for the formulated product.

While hazard assessment for most chemicals typically involves investigating the relationship between increasing exposure and increasing observed adverse effects in laboratory studies, some chemicals may have the potential to cause impacts at very low doses.

Examples of this are the endocrine disrupting chemicals (EDCs), which can interfere with an animal's endocrine (hormone) system, potentially at very low exposure levels. Certain chemicals such as the plasticizers found in plastic bottles are suspected to be EDCs. At the present time, there is no evidence that any of the herbicide active ingredients used in invasive weed control are EDCs. The US EPA studied glyphosate and 2,4-D through their [Endocrine Disruptor Screening Program](#) and determined that no

convincing evidence exists that either substance disrupts estrogen, androgen, or thyroid pathways. Studies have not been conducted for the other herbicides discussed in this manual, but none are on the European Union list of suspected endocrine disruptors.

Hazard Quotients Defined

The Hazard Quotient (HQ) is a measure of risk and is defined as the ratio of the predicted exposure to a Toxicity Reference Value (TRV) for the particular type of wildlife being assessed. HQ values >1 indicate that exposure exceeds the "No Effect" level, and wildlife may be at risk of adverse effects. For these exposure scenarios, action should be taken by the land manager to reduce exposure.

Table 10. Half-Maximum Application Rates Used in Risk Charts

Herbicide Active Ingredient	Half-Max Application Rate (lbs AE or AI per acre)	Half-Max Application Rate (rate per acre)
Aminopyralid	0.055	3.5 oz of Milestone®/acre
Chlorsulfuron	0.061	1.5 oz of Telar®/acre
Clopyralid	0.125	0.335 pints Transline®/acre
Glyphosate	4.0	3.5 quarts RoundupProMax®/acre (with surfactant) 4 quarts Aquamaster®/acre (no surfactant)
Imazapyr	0.75	3 pts Habitat®/acre
Triclopyr BEE	4.0	4 quarts Garlon 4®/acre
Triclopyr TEA	4.5	1.5 gals Garlon 3®/acre
2,4-D	2.0	4 pts Weedar®/acre


AE = Acid Equivalent; AI = Active Ingredient.

*Fluazifop (Fusilade®) is the one widely-used active ingredient not included in the risk charts because USFS risk analysis was completed after the risk charts were developed

How to Read the Risk Charts

In the risk charts that follow, risk is expressed as a Hazard Quotient (HQ), which is the ratio of the predicted exposure to a Toxicity Reference Value (TRV), a level of exposure that is anticipated to be without adverse effects.

Each bar on the chart shows a range of estimated risk for a specific exposure scenario based on three estimates of exposure—best-case (low exposure), most-probable (the most likely exposure), and worst-case (high exposure). Each estimate is based on a set of assumptions, such as the amount of herbicide residue on food (such as foliage, fruits, and insects) and the amount of food eaten or the amount of runoff into a water body. Factors used to estimate exposure specific to each scenario are listed in the caption for each chart.



The **best-case risk estimate** is at the left end of each bar and assumes the lowest exposure. The **most-probable risk estimate** (HQ=0.40 in the example above) is located at the point at which the bar changes color from light gray to dark gray and assumes the most likely exposure. The **worst-case risk estimate** is at the right end of the bar and assumes worst-case exposures.

The background of each risk chart is color-coded, with a HQ in the green zone indicating low risk, an HQ in the yellow zone indicating that anticipated exposures are approaching a level of concern, and an HQ in the red zone indicating that the predicted exposure will exceed the TRV, and adverse effects may result. Because wildlife TRVs are derived from No Observable Adverse Effect Levels (NOAELs), a bar in the red zone does not necessarily mean that harm will occur, but risks that fall in this zone should prompt the land manager to consider steps to mitigate the risk. The further the bar is into the red zone, the more likely it is that adverse effects will occur. The BMPs in Section 3 describe steps that can be taken to reduce risks when HQ values risk calculations exceed a level of concern.

The scale of the charts is logarithmic, which allows for the display of values that differ by many factors of ten. The logarithmic scale also visually compresses the bars and skews plots slightly to the right—for example, a HQ value of 0.5 is not exactly in the middle between 0.1 and 1, but slightly to the right of the halfway point.


Overview of Risks to Wildlife from Use of Common Herbicides

Overall, the risk estimates shown in the charts demonstrate that for the majority of the most-probable acute exposure scenarios, the herbicides pose low risks to wildlife. An exception to this involves fish and aquatic invertebrates exposed to glyphosate formulations that contain certain higher-toxicity surfactants such as polyethylenamine (POEA). These products cannot be legally applied directly to water, and applicators should also use caution when making applications near aquatic sites, such as ephemeral pools that may be used as breeding areas for amphibians and insects. Using glyphosate products that do not contain POEA in these settings can reduce the potential for impacts.

A second example of risks that may exceed the level of concern under the most-probable exposure scenarios involves products that contain either triclopyr BEE or triclopyr TEA. In these cases, the HQ values can exceed the level of concern for chronic exposure scenarios when large, herbivorous mammals consume vegetation that contains residues of these herbicides.

With regard to the worst-case (highest) exposure level scenarios, 2,4-D acid, glyphosate/surfactant combinations and triclopyr BEE and TEA can all pose risks that exceed the level of concern. These scenarios include both acute and chronic exposures for aquatic invertebrates, fish, mammals and birds.

Figure 4:



Taxa: Adult stage honey bees are used as a surrogate for all terrestrial insects.

Assumptions: Terrestrial application of herbicide at half of the maximum rate on a representative product's label (see Table 11); 50% of the bee's body surface is covered with herbicide; 100% of herbicide is absorbed; the distance between the bee and the sprayer is 0-10 feet.

Likelihood: Most likely with spray-to-wet applications on blooming plants or those with extrafloral nectaries.


Mitigation: Do not apply to blooming plants. Apply early in the morning or close to sunset when insects are less active. Use low-volume applications and reduce the amount applied per acre.

Risk calculated as a function of: The inherent toxicity of the herbicide to honey bees; the amount of active ingredient sprayed; and the distance between bee and applicator. Risks in this chart do not account for potential toxicity of any surfactants that are part of the product formulation or added to spray mixtures.

Methodology and sources: See description following risk charts and [PRI website](#), where you can access a spreadsheet for adjusting application rates and other variables.

Reading the chart: For each bar, the labeled central value is the most likely estimate. The right end of the bar assumes worst-case conditions for all underlying variables; the left end of the bar assumes best-case conditions. Mitigation is advised if risk enters the red zone.

Figure 5:



Taxa: Aquatic invertebrates.

Assumptions: Terrestrial application of herbicide at half of the maximum rate on a representative product's label (see Table 11); 10-acre treatment with no buffer zone between treatment area and water body.

Likelihood: Buffer zones may be required on some water ways and are common practice when using herbicides not approved for aquatic use. Dry season applications can result in long intervals before a rain event, resulting in lower residues for runoff.


Mitigation: Use low-volume applications and reduce the amount applied per acre. Use buffer zones (see Bakke (2001) to help gauge effective buffer distances). Make applications during the dry season to avoid runoff. For applications near waterways, consider using herbicide formulations intended for use in aquatic systems.

Risk calculated as a function of: The inherent acute toxicity of the herbicide to aquatic invertebrates; herbicide characteristics that affect transport through soil to water (water solubility, ability to adsorb to soil); soil type; and the application rate. Herbicide degradation is not considered, as the estimate is for runoff occurring soon after the application. Except for glyphosate with the POEA surfactant, risks in this chart do not account for potential toxicity of any surfactants that are part of the product formulation or added to spray mixtures.

Methodology and sources: See description following risk charts and [PRI website](#) where you can access a spreadsheet for adjusting application rates and other variables.

Reading the chart: For each bar, the labeled central value is the most likely estimate. The right end of the bar assumes worst-case conditions for all underlying variables; the left end of the bar assumes best-case conditions. Mitigation is advised if risk enters the red zone.

Figure 6:



*Exposure to TCP, the degradation product of triclopyr TEA and triclopyr BEE, is not reflected in the risk charts because first-flush runoff has been assumed to occur soon after application, before significant amounts of TCP have formed.

Taxa: Fish are also used as a surrogate for amphibians.

Assumptions: Terrestrial application of herbicide at half of the maximum rate on a representative product's label (see Table 11); 10-acre treatment with no buffer zone between treatment area and water body; rain within 24 hours of application.

Likelihood: Buffer zones may be required on many water ways and are common practice when using herbicides not approved for aquatic use. Dry season applications in California will result in a long interval before a rain event, resulting in lower residues for runoff.


Mitigation: Use low-volume applications and reduce the amount applied per acre. Use buffer zones (see Bakke (2001) to help gauge effective buffer distances). Make applications during the dry season to avoid runoff. For applications near waterways, consider using herbicide formulations intended for use in aquatic systems.

Risk calculated as a function of: The inherent acute toxicity of the herbicide to fish; herbicide characteristics that affect transport through soil to water (water solubility, ability to adsorb to soil); soil type; and the application rate. Herbicide degradation is not considered, as the estimate is for runoff occurring soon after the application. Except for glyphosate with the POEA surfactant, risks in this chart do not account for potential toxicity of any surfactants that are part of the product formulation or added to spray mixtures.

Methodology and sources: See description following risk charts and go to [PRI website](#) where you can access a spreadsheet for adjusting application rates and other variables.

Reading the chart: For each bar, the labeled central value is the most likely estimate. The right end of the bar assumes worst-case conditions for all underlying variables; the left end of the bar assumes best-case conditions. Mitigation is advised if risk enters the red zone.

Figure 7:



Taxa: Small mammals.

Assumptions: Terrestrial application of herbicide at half of the maximum rate on a representative product's label (see Table 11); 10-100% of diet is contaminated.

Likelihood: Under spot applications it is possible that a significant portion of a small mammal's diet could be contaminated. With broadcast applications over any sizable area (unusual for wildland management) contamination is likely for some small mammals.


Mitigation: Use low-volume application and reduce the amount applied per acre. If possible, don't treat large contiguous areas all at once. Avoid contamination of plants used as food sources by small mammals.

Risk calculated as a function of: The inherent acute toxicity of the herbicide to mammals; the residue rate of herbicide on fruit (which is proportional to the application rate). Except for glyphosate with the POEA surfactant, risks in this chart do not account for potential toxicity of any surfactants that are part of the product formulation or added to spray mixtures.

Methodology and sources: See description following risk charts and go to [PRI website](#), where you can access a spreadsheet for adjusting application rates and other variables.

Reading the chart: For each bar, the labeled central value is the most likely estimate. The right end of the bar assumes worst-case conditions for all underlying variables; the left end of the bar assumes best-case conditions. Mitigation is advised if risk enters the red zone.

Figure 8:



Taxa: Small mammals.

Assumptions: Terrestrial application of herbicide at half of the maximum rate on a representative product's label (see Table 11); 10-100% of diet is contaminated.

Likelihood: Under spot applications it is unlikely that a significant portion of a small mammal's insect-based diet could be contaminated. With broadcast applications over any sizable area (unusual for wildland management) contamination is possible for some small mammals.


Mitigation: Use low-volume applications and reduce the amount applied per acre. If possible, don't treat large contiguous areas all at once. Avoid treating plants when feeding by insects is likely, if known.

Risk calculated as a function of: The inherent acute toxicity of the herbicide to mammals; the residue rate of herbicide on insects (which is proportional to the application rate). Except for glyphosate with the POEA surfactant, risks in this chart do not account for potential toxicity of any surfactants that are part of the product formulation or added to spray mixtures.

Methodology and sources: See description following risk charts and go to [PRI website](#), where you can access a spreadsheet for adjusting application rates and other variables.

Reading the chart: For each bar, the labeled central value is the most likely estimate. The right end of the bar assumes worst-case conditions for all underlying variables; the left end of the bar assumes best-case conditions. Mitigation is advised if risk enters the red zone.

Figure 9:



Taxa: Large mammals.

Assumptions: Terrestrial application of herbicide at half of the maximum rate on a representative product's label (see Table 11); 10-100% of diet is contaminated for several months.

Likelihood: Under spot applications it is unlikely that a significant portion of any large mammal's diet would be contaminated. With broadcast applications over any sizable area (unusual for wildland management) consider the feeding range of the wildlife relative to the treatment area.

Mitigation: Use low-volume applications and reduce the amount applied per acre. If possible, don't treat large contiguous areas all at once. Avoid contamination of plants known to be used as food sources by large mammals.

Risk calculated as a function of: The inherent chronic toxicity of the herbicide to mammals; the residue rate of herbicide on vegetation (proportional to the application rate). Except for glyphosate with the POEA surfactant, risks in this chart do not account for potential toxicity of any surfactants that are part of the product formulation or added to spray mixtures.


Methodology and sources: See description following risk charts and go to [PRI website](#), where you can access a spreadsheet for adjusting application rates and other variables.

Reading the chart: For each bar, the labeled central value is the most likely estimate. The right end of the bar assumes worst-case conditions for all underlying variables; the left end of the bar assumes best-case conditions. Mitigation is advised if risk enters the red zone.

Figure 10:



Chronic Risks to Large Birds from Consuming Contaminated Vegetation



Taxa: Large birds.

Assumptions: Terrestrial application of herbicide at half of the maximum rate on a representative product's label (see Table 11); 10-100% of diet is contaminated for several months.

Likelihood: Under spot applications it is unlikely that a high portion of any bird's diet would be contaminated. With broadcast applications over any sizable area (unusual for wildland management) consider the feeding range of the wildlife relative to the treatment area.


Mitigation: Use low-volume applications and reduce the amount applied per acre. If possible, don't treat large contiguous areas all at once. Avoid contamination of plants known to be used as food sources by birds. Avoid treatments during nesting season.

Risk calculated as a function of: The inherent chronic toxicity of the herbicide to birds; the residue rate of herbicide on vegetation (which is proportional to the application rate). Except for glyphosate with the POEA surfactant, risks in this chart do not account for potential toxicity of any surfactants that are part of the product formulation or added to spray mixtures.

Methodology and sources: See description following risk charts and go to [PRI website](#), where you can access a spreadsheet for adjusting application rates and other variables.

Reading the chart: For each bar, the labeled central value is the most likely estimate. The right end of the bar assumes worst-case conditions for all underlying variables; the left end of the bar assumes best-case conditions. Mitigation is advised if risk enters the red zone.

Figure 11:



Taxa: Small birds.

Assumptions: Terrestrial application of herbicide at half of the maximum rate on a representative product's label (see Table 11); 10-100% of diet is contaminated.

Likelihood: Under spot applications it is unlikely that a high portion of any bird's insect-based diet would be contaminated. With broadcast applications over any sizable area (unusual for wildland management) consider the feeding range of the wildlife relative to the treatment area.

Mitigation: Use low-volume applications and reduce the amount applied per acre. If possible, don't treat large contiguous areas all at once. Avoid treating plants when insects are feeding. Avoid treatments during nesting season.

Risk calculated as a function of: The inherent acute toxicity of the herbicide to birds; the residue rate of herbicide on insects (which is proportional to the application rate). Except for glyphosate with the POEA surfactant, risks in this chart do not account for potential toxicity of any surfactants that are part of the product formulation or added to spray mixtures.

Methodology and sources: See description following risk charts and go to [PRI website](#), where you can access a spreadsheet for adjusting application rates and other variables.

Reading the chart: For each bar, the labeled central value is the most likely estimate. The right end of the bar assumes worst-case conditions for all underlying variables; the left end of the bar assumes best-case conditions. Mitigation is advised if risk enters the red zone.

Risk Assessment Methodology

The methods used for estimating risk are based closely on USFS risk assessment methodology ([link](#)), in which three estimates are calculated for the exposure (dose) received as a result of various herbicide use scenarios. Each dose estimate is based on a set of best-case, most-probable, or worst-case assumptions based on exposure parameters appropriate to that scenario. The dose estimates are then compared to Toxicity Reference Values to assess risk if the scenario were to occur.

Exposure estimates were calculated using the risk assessment spreadsheets developed by Syracuse Environmental Research Associates (SERA) for the USFS and the Bureau of Land Management (BLM), published between 2007 and 2014. A full description is available in the report "[Preparation of Environmental Documentation and Risk Assessments](#)." Risk assessments for each of the herbicides discussed here are also downloadable from the [USFS site](#). A detailed explanation of the methods used to estimate risk in this report is also available in Chapter 2 of the "[2010 Marin Municipal Wastewater District \(MMWD\) Herbicide Risk Assessment](#)." However, some parameter values and methods used for the risk estimates above differ from the *2010 MMWD Herbicide Risk Assessment*. Each of these changes is discussed below. Finally, the [PRI website](#) provides detailed information on how the risk charts were developed and allows users to modify application rates to assess changes in risk profiles.

Modifications to USFS Risk Estimation Methods
Several modifications to USFS/SERA default values were made for this evaluation:

TRVs: Toxicity Reference Values (TRVs) based on LD₅₀ or LC₅₀ transformed to "No Effect" levels by incorporating an additional uncertainty factor of 20, the methodology used by US EPA to adjust TRVs for assessment effects to endangered species. This transformation ensures that all TRVs are based on "No Effect" levels and allows direct comparison of herbicides. This change has been incorporated into the more recent USFS herbicide risk assessments, and PRI updated the older risk assessments to include this change.

Percent of diet contaminated: In more recent versions of the USFS/SERA herbicide risk assessments, the percentage of an animal's diet assumed to be contaminated was modified to 10% (best-case), 30% (most-probable) or

100% (worst-case). PRI applied the same change to herbicides not yet adopted by USFS, to ensure an "apples to apples" comparison between herbicides. Residue rates assumed for herbicides on food (fruit, vegetation and prey) were based on the most up-to-date values from USFS/SERA (WorksheetMaker 6.0). The caloric error factor, which was introduced in recent versions of USFS/SERA worksheets, was not utilized here.

Herbicide Residue Rates: USFS changed the residue rates used in the latest version of their risk calculation spreadsheets for estimating exposures from consumption of contaminated fruit, insects and vegetation. This change lowers the best-case predicted dose for wildlife from consumption of contaminated food. In the new versions of the spreadsheets, a new lower residue rate was introduced that is equivalent to the following:

Best-case residue rate = Most-probable rate x (Most-probable rate ÷ Worst-case rate)

These values were incorporated into the calculations for all of the herbicides to ensure comparison of equivalent value.

Insect Contamination Rate: The USFS changed the mass of a honey bee from 93 mg to 116 mg and the surface area from 2.66 cm² to 1.42 cm² in the more recent herbicide reviews. The net effect is to reduce the estimated dose received by the honey bee. These values were incorporated into the calculations for all of the herbicides to ensure comparison of equivalent values.

Toxicity Reference Values Used to Estimate Risk

Toxicity Reference Values (TRVs) are given in terms of mg of acid equivalent (AE) or active ingredient (AI).
NOAEL is the No-Observed-Adverse-Effect Level.

Table 11. Toxicity reference values used to estimate risk

RECEPTOR (UNITS)	HERBICIDE	TRV USED	USFS TRV	ENDPOINT
Honeybees (mg/bee)	2,4-D Acid	1075	1075	NOAEL
	Aminopyralid	1075	1075	NOAEL
	Chlorsulfuron	25	25	NOAEL
	Clopyralid	909	909	NOAEL
	Glyphosate	860	860	NOEC
	Imazapyr	860	860	NOAEL
	Triclopyr BEE	620	620	NOAEL ^b
	Triclopyr TEA	620	620	NOAEL ^b
Birds, acute (mg/kg body weight)	2,4-D Acid	415	415	NOAEL
	Aminopyralid	14	14	NOAEL
	Chlorsulfuron	1686	1686	NOAEL
	Clopyralid	670	670	NOAEL
	Glyphosate	1500	1500	NOAEL
	Imazapyr	2510	2510	NOAEL
	Triclopyr BEE	126	126	NOAEL ^b
	Triclopyr TEA	126	126	NOAEL ^b
Birds, chronic (mg/kg body weight)	2,4-D Acid	76	76	NOAEL
	Aminopyralid	184	184	NOAEL
	Chlorsulfuron	140	140	NOAEL
	Clopyralid	15	15	NOAEL
	Glyphosate (no surfactants)	58	58	NOAEL
	Glyphosate (with POEA)	43	43	NOAEL
	Imazapyr	610	610	NOAEL
	TCP ^c	116	116	NOAEL ^b
	Triclopyr BEE	7.5	7.5	NOAEL ^b
	Triclopyr TEA	7.5	7.5	NOAEL ^b
Mammals, small (mg/kg body weight)	2,4-D Acid	25	25	NOAEL
	Aminopyralid	104	104	NOAEL
	Chlorsulfuron	75	75	NOAEL
	Clopyralid	75	75	NOAEL
	Glyphosate	500	500	NOAEL
	Imazapyr	738	738	NOAEL
	Triclopyr BEE	440	440	NOAEL ^b
	Triclopyr TEA	440	440	NOAEL ^b

Receptor (units)	Herbicide	TRV Used	USFS	Endpoint
Mammals, large (mg/kg body weight)	2,4-D Acid	5	5	NOAEL
	Aminopyralid	50	50	NOAEL
	Chlorsulfuron	5	5	NOAEL
	Clopyralid	15	15	NOAEL
	Glyphosate	500	500	NOAEL
	Imazapyr	738	738	NOAEL
	TCP ^c	12	12	NOAEL ^b
	Triclopyr BEE	0.4	0.4	NOAEL ^b
	Triclopyr TEA	0.4	0.4	NOAEL ^b
Fish (mg/liter of water)	2,4-D Acid	4.8	95.6	LC ₅₀ ÷ 20
	Aminopyralid	50	50	NOEC
	Chlorsulfuron	30	30	NOEC
	Clopyralid	5 ^a	103	LC ₅₀ ÷ 20
	Glyphosate (no surfactants)	0.5	0.5	NOAEC
	Glyphosate (with POEA)	0.048	0.048	NOAEC
	Imazapyr	10.4	10.4	NOAEC
	TCP ^c	0.18	0.18	NOAEC ^b
	Triclopyr BEE	0.091	0.091	NOAEC ^b
	Triclopyr TEA	20	20	NOAEC ^b
Aquatic Invertebrates (mg/liter of water)	2,4-D Acid	1.25 ^a	25	LC ₅₀ ÷ 20
	Aminopyralid	89	89	NOEC
	Chlorsulfuron	10	10	NOEC
	Clopyralid	23.1	23.1	NOEC
	Glyphosate (no surfactants)	2.7	2.7	NOAEC
	Glyphosate (with POEA)	0.075	0.075	NOAEC
	Imazapyr	41	41	NOAEC
	TCP ^c	0.55	0.55	NOAEC ^b
	Triclopyr BEE	0.045	0.045	NOAEC ^b
	Triclopyr TEA	25	25	NOAEC ^b

^A To ensure comparison of equivalent endpoints between herbicides, all TRVs values expressed as LC₅₀ or LD₅₀ values were translated by either USFS or PRI to “No Effect” levels by incorporation of an uncertainty factor of 20, similar to that used by US EPA to protect endangered species. This practice was only recently incorporated into the USFS methodology, so PRI implemented these changes for the herbicides reviewed by USFS prior to the change.

^b For triclopyr and TCP toxicity to mammals, USFS used allometric parameters that correct the NOAEL for the amount of food and water consumed, based on body weight and size, to adjust for differences between the test species and the taxa to which the TRV is applied.

^c TCP is the primary degradation product of triclopyr. Because triclopyr must degrade before any TCP is produced, only the chronic scenarios of large mammals and birds eating vegetation involve potential exposure to TCP. The other scenarios are acute events, where triclopyr has not yet degraded to form TCP. Chronic exposure to treated vegetation will result in exposure to a combination of the parent compound and TCP, which degrade at similar rates. The risk bars are based on the TRV for the more toxic (lower value) of the two to produce a more protective risk estimate. For both mammals and birds, the risk charts are based on the TRV for triclopyr acid, since it has the lower value.

Factors Affecting Herbicide Runoff to Surface Waters

Herbicide Half-Life

Herbicide half-life is a measure of persistence in the environment. Herbicides that are persistent in the soil environment continue to have herbicidal activity and cause adverse effects on the ecosystem until the concentration drops below a level that is toxic to plants. The range of half-lives for the herbicides in soil under aerobic conditions—in the presence of oxygen and microbes—can vary by a factor of ten or more for each herbicide. Exposure to sunlight can accelerate decomposition of some herbicides. The longest half-lives are typically relevant under arid conditions where microbial degradation rates are low. Anaerobic degradation is usually slower than aerobic degradation. In general, glyphosate is expected to be less persistent than other herbicides considered in this assessment, while imazapyr and aminopyralid are among the most persistent. Triclopyr BEE and TEA rapidly degrade or dissociate to triclopyr acid, so the persistence of triclopyr degradates—triclopyr acid and TCP—is most relevant to triclopyr applications. Organic herbicides such as clove oil, pelargonic acid, and limonene have very short half-lives (a few days to a week), which limits their potential for exposure.

Figure 4 shows the range of half-lives for the herbicides in soil under aerobic conditions. In the plot, herbicides are arranged in order of the Central value of their measured half-life. The Upper, Lower and Central values on Figure D-1 are based on a review of the academic literature and the values used by government agencies, including US EPA, USFS, California Department of Pesticide Regulation (DPR), and the Oregon Department of Environmental Quality (ODEQ) (see [PRI website](#) for more information).

The Central values for the herbicides used in the plots (except for 2,4-D and aminopyralid) in Figure D-1 are the half-life values used by USFS in its risk assessments as the Central half-life estimate in soil, with the values for 2,4-D from DPR's environmental fate review and for aminopyralid from US EPA's risk assessment. Lower and Upper values used in the figure are taken from US EPA's risk assessments or from DPR's or ODEQ's environmental fate documents summarizing the available literature studies. Half-lives vary depending on test conditions, and comparable studies conducted under the same test conditions were not always available for every herbicide. When soil values were unavailable, the half-life on fruit was used.

Figure 4 is intended to provide as much as possible an “apples-to-apples” comparison of aerobic soil half-lives. However, imazapyr does not degrade in soil under aerobic conditions, so a field dissipation half-life (5.9 years) is used, in order to provide a numerical point of comparison to other herbicides. Note that half-lives of herbicides in water or in anaerobic sediments (such as wetlands) may be different than the aerobic soil half-lives presented in Figure 4. For most pesticides, the anaerobic half-life (in the absence of oxygen) is longer than the aerobic half-life. Sunlight and processes that dissipate herbicides in the environment like rainfall runoff, absorption by plants, or irreversible binding to soils can also alter the persistence of a chemical in the treated area.

Figure 4 shows the total range of half-lives observed for the different chemicals. Half-life values used by the USFS in their worksheets are those used to produce the charts and are more narrowly constrained to reflect half-lives under the most common conditions.




Figure 12: Comparison of the range of herbicide half-lives under aerobic conditions in soil. The high end of the range is typically under arid conditions where microbial degradation rates are low. Exposure to sunlight can accelerate decomposition and shorten the half-life of some herbicides. Sources are described after Table 12. For aminopyralid, see [EPA Fact Sheet 2005](#). For imazapyr, see [EPA 2007 Appendix A Imazapyr Effects Determination for the CA Red-legged frog](#).

Water Contamination Rates

Water contamination rates are a measure of how much of an applied herbicide will run off of the treated area into nearby water bodies. Maximum or peak concentrations of herbicides in water bodies receiving runoff are typically observed when rainfall or irrigation occurs soon after treatment, before the herbicide has degraded substantially. The concentration of herbicide in this “first-flush” runoff may potentially impact aquatic organisms and terrestrial animals that make contact with or drink contaminated water. The potential of herbicides to move off-site in runoff water depends on water solubility, half-life, and the ability of the herbicide to bind to soil. The site characteristics are relevant too, as different soil types bind to herbicides differently. Bare or impermeable soils are much more prone to runoff than vegetated areas; sandy soils are susceptible to leaching that may result in groundwater contamination.

The risk charts use the USFS method (based on the Groundwater Loading Effects of Agricultural Management Systems (GLEAMS) model) to estimate the concentration of each herbicide in water for an application to 10 acres, no buffers along the edge of the treated area, and rainfall after the application based on averages for a variety of sites. The range of water contamination rates is based on the range of site variables such as soil type and chemical properties. Use of buffer zones around water bodies will reduce water contamination.

Water contamination rates are measured in units of milligrams of herbicide per liter per pound of herbicide applied per acre (mg/L per lb/acre). Actual herbicide concentrations in the receiving water body will

depend on how many pounds of active ingredient are applied to land that drains to the water body. Use of herbicides with application rates of fractions of a pound per acre (see Table 11) will generally result in lower concentrations than herbicides with higher application rates. Predicted concentrations in the receiving water bodies for the half-maximum application rates for each active ingredient are shown in Figure 5. These concentrations were used to estimate the risks displayed in the charts for aquatic species and for animals drinking the water.




Figure 13: Comparison of the range of predicted concentrations in peak runoff after terrestrial application at half-maximum application rate. Factors affecting predicted concentrations include application rate, water solubility, half-life, and the ability of the herbicide to bind to soil (Koc). Use of buffer zones near surface waters will help to reduce water contamination. Source: “Estimated Water Contamination Rates” in USFS risk assessment worksheets at www.fs.fed.us/foresthealth/pesticide/worksheets.shtml.

Herbicide Selection Process

Several factors contributed to selecting herbicides to control noxious weeds on Clallam County roadsides.

- **EPA Approved for Roadside Use** -- All the selected herbicides are fully labeled for use on roadsides and are registered for use in the state of Washington. The herbicide label does not have to list all the weeds, but the label does have to list roadsides or rights-of-way as a use site.
- **Effectiveness on Target Species** – AquaNeat and Polaris are very broad spectrum and will control most of the county's noxious weeds. Milestone, Transline, Vastlan, and 2,4-D are selective and very effective only on broadleaf plants. Fusillade II is effective only on grass species. Many of the targeted weeds have a perennial life cycle with persistent root systems. Effective control requires translocated herbicides that kill the roots. All the selected herbicides translocate to the roots.
- **Selectivity** – Several of the herbicides were chosen because they selectively target broadleaf weeds, not grasses. This allows grass to be unaffected and to colonize space previously occupied by broadleaf weeds. The grass herbicide gives the program a selective chemical for controlling weedy grasses, such as reed canarygrass, in a mixed plant community.
- **Human and Environmental Safety** – A carefully selected, limited palette of herbicides rated with low toxicity to humans and wildlife has been chosen for this program. Some of the products are labeled for aquatic use so inadvertent occurrences in water is anticipated to have minimal effects on aquatic organisms. Most are labeled for use on grazed areas such as range and pasture. Most are labeled for use in maintaining wildlife habitat, fence rows, as well as rights-of-way.

Herbicide Product List

Clallam County proposes to use the following products for targeted herbicide applications:

AquaNeat® (aquatic formulation glyphosate)
Element 3A® (aquatic formulation triclopyr)
Fusillade II® (fluazifop-P)
Garlon 3A® (aquatic formulation triclopyr)
Garlon 4® (triclopyr)
HighNoon® (aquatic formulation aminopyralid and florpyrauxifen)
Milestone® (aquatic formulation aminopyralid)
Oust XP® (sulfometuron-methyl)
Plateau® (imazapic)
Polaris® (aquatic formulation imazapyr)
Ranger Pro® (glyphosate)
Transline® (clopyralid)
Vastlan® (aquatic formulation triclopyr)
WeeDestroy AM-40® (aquatic formulation 2,4-D)

The chosen products are effective on known roadside and pit weeds, offer the greatest weed selectivity, maximize worker and public safety (no wait, access when the spray has dried), and used as proposed, pose low risk for wildlife and the environment. See Appendix B for risk analysis.

The standard, minimum personal protection equipment (PPE) when using herbicides includes:

Long sleeved shirt, long pants
Shoes plus socks
Chemical resistant gloves made of any waterproof materials

The required PPE for each herbicide, listed on the label, will be strictly adhered to during treatment.

Table 12. Selected herbicide characteristics.

Chemical Name Product Name	Selective	Aquatic Sites	Target Weeds	Personal Protection Equipment	Comments
<u>2,4-D</u> WeeDestroy AM-40	✓	✓	Broadleaf	Standard; eye protection + apron for mixing	Inexpensive, often used in mix; short residual
<u>Aminopyralid</u> ¹ Milestone	✓	✓	Broadleaf	Standard	Moderate residual may help suppress seed germination; very low rates
<u>Aminopyralid and florpyrauxifen</u> HighNoon	✓	✓	Broadleaf, some annual grasses	Standard	Moderate residual may help suppress seed germination; very low rates
<u>Clopyralid</u> Transline	✓		Broadleaf	Standard	Very selective; will not affect many native and desirable plants; moderate residual; low rates
<u>Fluazifop-P</u> Fusilade II	✓		Grasses	Standard + eye protection	For dry sites; reed canary-grass and annual grasses
<u>Glyphosate</u> AquaNeat		✓	All weeds	Standard	Minimal to no residual; protect desirable vegetation
<u>Glyphosate</u> Ranger Pro			All weeds	Standard	For pits and some special sites only
<u>Imazapic</u> Plateau	✓		Annual weeds	Standard	For dry sites, annual weeds, and for prairie restoration, wildflower establishment
<u>Imazapyr</u> Polaris		✓	All weeds	Standard	Long residual; protect desirable vegetation
<u>Sulfometuron- methyl</u> Oust XP			Broadleaf, grasses	Standard	Long residual; early season pre and post-emergent; for use in county pits only
<u>Triclopyr amine</u> Element 3A, Garlon 3A	✓	✓	Broadleaf, shrubs	Standard; Eye protection for mixing, access to eye wash station	Moderate residual
<u>Triclopyr choline</u> Vastlan	✓	✓	Broadleaf, shrubs	Standard	Moderate residual
<u>Triclopyr ester</u> Garlon 4	✓		Broadleaf, shrubs	Standard	Moderate residual

¹Registered as a reduced risk pesticide under the EPA reduced risk pesticide program

Adjuvants are compounds added into an herbicide mix to improve efficacy. They perform various functions, including enhanced plant uptake of the herbicide; better mixing of otherwise incompatible herbicides; increased adhesion of the spray to plant surfaces; and reduced spray drift. In many herbicide products, adjuvants are included as part of the pre-mixed formulation as purchased. Applicators can also add adjuvants to spray mixtures prior to application. Adjuvants include marker dyes, which are visible indicators of freshly treated weeds, including Blazon and Highlite (aquatic formulation).

Surfactants, or "surface active agents", are a type of adjuvant added to a mix to increase the dispersing, spreading, wetting, or other properties of the liquids. Surfactants disperse water droplets and help penetrate a plant's waxy surface. (Table 13).

Some states require registration of adjuvants as pesticide products, but the US EPA does not, so relatively less is known about adjuvants compared to pesticide active ingredients. Acute toxicity information is often available, with some of these compounds being labeled as strong eye or skin irritants, but information regarding chronic toxicity is sparse. Washington State and European countries require environmental toxicology data on adjuvants.

For many pesticide products containing adjuvants as part of the formulation, the compounds are not explicitly identified on the label or the Safety Data Sheet. Unless they are on one of US EPA's lists of more toxic chemicals, they do not have to be identified. The identity of these ingredients in a pesticide or adjuvant product is legally protected from full disclosure as "Confidential Business Information."

Without more detailed information, it is not possible to conduct a comprehensive risk assessment on adjuvants, so they are not included in the risk charts shown as part of Appendix B, which focus on herbicidal active ingredients. However, at least one adjuvant is known to pose hazards to wildlife—the surfactant used in the original formulation of RoundUp®, polyoxyethyleneamine (POEA). This surfactant is more toxic to aquatic life than the active ingredient glyphosate—it has been included as a separate entry in the risk charts. Nonylphenol ethoxylates (NPEs), which are used in some adjuvants (and many consumer products), may be linked to endocrine disrupting effects. No products containing polyethoxylated tallowamine (POEA) or nonylphenol ethoxylates (NPEs) will be allowed for use in this program. Adjuvants with low toxicity to wildlife include modified seed oils, alkyl ethoxylates, and silicones. Liberate®, Competitor®, DyneAmic®, Syl-Tac EA® and Agri-Dex® (all aquatic formulations) are brand names of some adjuvants from these low toxicity categories and have been selected for use in this program. Research is developing on this subject and will be regularly added to updates for this program.

Government agencies negotiate for favorable pricing and award a contract to a preferred provider for many goods and services. Herbicides will be purchased under state contract whenever possible to conserve tax dollars. Because the preferred provider may vary from year to year; different brand names than listed in the previous tables, with the same active ingredient may be substituted. New products or different formulations with the same active ingredient that are more user or environmental friendly and cost beneficial will be substituted as they become available.

Table 13. Adjuvants used to enhance herbicide effectiveness.

Adjuvants	Aquatic use	Treatment effects	PPE	Comments
Competitor - vegetable oil Agri-Dex, - crop oil concentrate Dyne-Amic - nonionic surfactant Liberate - fatty acids Hasten- vegetable oil Syl-Tac EA, organosilicone surfactant and modified vegetable oil concentrate	✓	Increases herbicide uptake	Standard	Used at low rates
Blazon - marker dye Highlite - marker dye	✓	No active effect	Standard	Highlights recently sprayed weeds; washable

Several studies have shown non-synthetic products (or “natural”) are considerably less effective for controlling weeds, especially biennials or perennials, than synthetic ones. However, three of these products, acetic acid, clove oil, and limonene are the subject of an on-going study for control of the annual weed, herb Robert. Pending study results, one or more of these herbicides may be added to the herbicide product list for control of this or other annual weeds.

Application Methods

Foliar. Applications to the plants' leaves are an easy way to control weeds with maximum amount of herbicide directed to the target plants and optimum up take by the plants for both herbaceous forbs and grasses.

Wiping Applicators. Wiping applicators (also called rope wicks) rub the concentrated herbicide solution on the plant's leaf and stem surfaces. Because only the weeds tall enough to contact the rubbing surface are affected, nonselective herbicides can be used selectively to release low-growing plants or plants below the treatment height. Drift does not occur with wiping applicators so there is no potential exposure for adjacent crops and gardens.

Stem Injection. Some species, such as knotweeds, have stems of sufficient size that herbicide can be injected directly into the stem. While this is an effective treatment, it is a very labor-intensive treatment for treating dense stands. Only some herbicides are labeled for this application method.

Stem Injection/Spaced Cuts/Cut Surface/Cut Stump/Basal Bark. Stem injection, spaced cuts, cut surface, cut stump, and basal bark are treatments often used for controlling tall growing woody plants. As the name implies, herbicide is applied to just the cut surface or the woody stem. The herbicide rate and carrier are adjusted according to the part of the woody plant being treated. Unlike foliar treatments done during the growing season, these treatments can be applied year-round. These treatments are particularly effective for large butterfly bush and Scotch broom in excess of 1-2 inches in diameter.

APPENDIX C: SAMPLE RECORD KEEPING FORMS

Treatment Form

2024 CLALLAM COUNTY ROADS : Herbicide/Manual Treatment Data Form

Name of Entity/Person for whom Treatment was applied: <u>Clallam County</u>			
Street Address: <u>223 E 4th St</u>		City: <u>Port Angeles</u>	State: <u>WA</u> Zip: <u>98362</u>
Address or Exact Location of Site: _____			
PIN#: _____			

General Activity Fields

County (circle one)	WRIA (circle one)	Project Name (from project list)	Department (circle one)	Workforce**
Clallam	15 16 17 18 19 20		Roads Parks DCD Other	

**Workforce: County Name, WCC Crew Name, County Weed Board

Crew Members Present:

Site/Inventory Fields

Date	Acres examined for weeds	Acres treated (do not lump plants)	Miles Examined	Miles Treated	Treatment Site (circle one)			Treatment Method (circle one)	Total <u>Manual</u> Infested Area Treated: (DO NOT lump plants together) acres
					Road Pit	Special Site Other	Private	Spot Complete Retreatment	

Weeds Treated (Just the PLANTS code is OK)	Infested Area Treated (DO NOT lump plants together)	% of area examined for weeds infested with species (<u>lump</u> plants together – use cover classes 1 - 9 listed below)	Manual/Herbicide	Number of plants manually removed	Notes

Cover Classes: 1 = Trace, 2 = 1 – 3%, 3 = 3 – 5%, 4 = 5 – 10%, 5 = 10 – 25%, 6 = 25 – 50%, 7 = 50 – 75%, 8 = 75 – 95%, 9 = 95 – 100%

Note: Cover classes are meant to be *approximations* only.

Herbicide ApplicationAll Licensed Applicators: Name and License # ☐ Christina St John #104740 ☐ Joe Oakes #100454 ☐ Sam Fischbein #102791Firm Name: Clallam County Noxious Weed Control Board Phone # 360-417-2442Firm Address: 223 E 4th St, Suite 15 City: Port Angeles State: WA Zip: 98362

Application Date	Time Start	Time Stop	Temp (F)	Wind Speed (MPH)	Wind Direction	Cloud Cover	Remarks - <u>Weather</u> forecast

Application Area (acre)	Total Volume of Mix Applied (gal)	Diluent	Special comment
		Water	

Product Name	EPA Registration #	Amount of herbicide used (oz)	Herbicide Applied/Acre or other measure	Concentration Applied
☐ Element 3A	62719-37			
☐ Milestone	62719-519			
☐ <u>Vastlan</u>	62719-687			
☐ Transline	62719-259			
☐ Competitor	WA-2935-04001			
☐ Blazon Blue				
☐ Oust XP	432-1552			
☐ <u>Aquaneat</u>	228-365			

Was this application made as a result of a permit? **Yes No**
If yes, Permit # _____Project Complete? **Y** or **N** (add notes)

Project ID #:

WA State NPDES Acres:

Notes:

Interactions:

Roadside Vegetation Monitoring Form**Clallam County
Weed Treatment Monitoring**

Examiner Name: _____

Evaluation Date: _____

Ref #	
Project # and Name	
<u>From "Comments":</u> Road name with BMP & EMP -OR- Min and Max Address	
Date(s) of treatment	
Herbicide or Manual treatment (circle one)	

Weeds Treated (Scientific name or code)	Infested Area Treated (acres)	Cover class from "% area examined for weeds infested with this species"	Percent efficacy of treatment (use codes on next page)

Do you think this treatment area is a high priority for retreatment next year? Yes / No

Please provide comments on the next page, if you have any.

Instructions: All information on page 1 of this datasheet comes from the “Herbicide/Manual Treatment Data Form”, except for:

- Examiner name
- Evaluation Date
- Percent efficacy of treatment

For Percent efficacy of treatment, enter the code that best approximates the percent of the population that was eradicated:

Code	% Efficacy	Rating	Description
0	0	No effect	No effect can be detected on the target species population
03	1 – 5	Failure	Little to no effect can be detected on the target species population.
15	6 – 25	Poor	Treatment killed less than a quarter of the target species population.
35	26 – 50	Marginal	Less than half of the target species population was controlled.
65	51 – 75	Fair	Over half of the target species population was controlled.
85	76 – 90	Good	Treatment was successful in killing most of the target species population
95	91 – 99	Excellent	Over 95% of the target species population has been killed with the treatment.
100	100	Complete	Not a single individual of the target species population was found after a complete survey of the site. The infestation was eradicated.
UN	UNK	Unknown	Treatment efficacy/success cannot be determined.

Comments:

APPENDIX D: SAMPLE PRESS RELEASE AND PUBLIC NOTICE

PUBLIC NOTICE

Clallam County is beginning the 20__ Integrated Weed Control program which may include spot treatments of herbicide to control specific noxious weeds and invasive species of special concern along selected portions of county right-of-way. Notices indicating which herbicide has been applied, the application date, and the target weed species will be posted onsite. The Integrated Weed Management Plan, which contains information about target weeds, locations, and treatment methods, can be viewed online at <https://www.clallamcountywa.gov/821/Noxious-Weed-Control>. Property owners who do not wish to have their adjoining right-of-way treated with herbicide have the option of keeping the right-of-way abutting their property weed free by applying for an Owner Will Control Agreement with Clallam County available online. Contact the County for further information at 360-417-2442.

NOTICE

The herbicides aminopyralid, imazapyr, triclopyr, clopyralid or _____ will be applied to this site to control noxious weeds, which threaten native vegetation and habitat in this area.

Planned / Actual Application Date*: _____

*Actual date of application contingent upon weather conditions.

Targeted Noxious Species:** _____

**Other weed species in this area may also be treated at this time.

NO USE RESTRICTIONS ARE IN PLACE

Avoid contact with treated vegetation until after it has dried.

Clallam County Noxious Weed Control Board
223 East Fourth Street, Suite 15
Port Angeles, WA 98362
(360) 417-2476
(360) 477-1210

APPENDIX F: OWNER WILL CONTROL PACKET

Available online at: <https://www.clallamcountywa.gov/821/Noxious-Weed-Control>



OWNER WILL CONTROL APPLICATION

Introduction:

The Noxious Weed Control Board thanks you for your interest regarding the Clallam County Integrated Weed Management Plan and the Owner Will Control opportunity. This packet contains an Owner Will Control Agreement application to be filled out completely and returned to the Noxious Weed board by mail or email. The Agreement releases all noxious weed control responsibility to the agreement holder dependent on Owner's compliance with the County control guidelines.

Owner Will Control Agreement:

The Agreement application must be completely filled out and returned to the Noxious Weed Board at:

Clallam County Noxious Weed Control Board (Or) **web_weed@clallamcountywa.gov**
223 E 4th St. Suite 15
Port Angeles, WA 98362

Incomplete applications cannot be accepted. The County will return the agreement holder a copy of the completed application at the email or mailing address provided within ten (10) business days of receipt.

It is important to return the application as soon as possible to ensure the county work plan can be modified to honor the application request. The County will email a reminder notice regarding applications for the following year to all current agreement holders.

Control Guidelines:

Appendix F.1 of this document includes best management practices for Owner control methods. The Owner must adhere to the county guidelines for effective control for each designated species found on county right-of-way. Failure to completely remove weeds as stipulated by this agreement will prompt a ten (10) day notice to comply with control guidelines or the Agreement may be terminated.

Native Plant Enhancement Program:

The goal of this program is to create natural, site appropriate plant communities along county road sides. Once weeds are controlled Owners may be eligible for native material and pollinator friendly plants from the County. Please indicate your interest in this program on your application.



OWNER WILL CONTROL AGREEMENT

By entering into this agreement an adjacent property owner (hereinafter referred to as "Owner") will agree to control noxious weeds and other weeds of concern as described in Appendix A of this agreement on county right-of-way adjacent to property located at:

(Street)

(City)

(Zip)

The County will send a confirmation email upon receiving a completed application and return a copy of the finalized Owner Will Control Agreement (hereinafter referred to as "Agreement").

For the purpose of this Agreement, 'control' will consist of complete removal of all above ground biomass and as much of the root system as is feasible of weeds listed in your packet, as well as any additional weeds of concern as determined by the County.

If noxious or other weeds of concern are observed on right-of-way adjacent to above named address, County will notify property owner of their presence. Owner will then have ten (10) days to completely remove weeds as required by this Agreement. If Owner fails to control weeds in that timeframe, this Agreement will be terminated and weeds will be controlled as determined by the County, including the use of herbicides.

This Agreement is valid from the date signed by both parties until December 31 of the same year.

If the Owner Will Control Agreement is terminated as described above the Owner may apply to re-enter into a new Owner Will Control Agreement the following calendar year.

* _____ * _____ * _____

Owner Name (Print)

(Signature)

Date

* _____ * _____

(Owner Email)

(Owner Phone #)

Interested in Native Plant Enhancement Program?

(circle one)

YES

NO

* _____ * _____ * _____

County Representative

(Signature)

Date

*Required Field



Appendix F.1:

Owner will be responsible for the complete control of species listed below in the accordance to the county guidelines.

Species List for Control:

<i>Vegetative Propagation*</i>		<i>Non-Vegetative Propagation</i>	
bindweed, field	laurel, spurge	alyssum, hoary	herb Robert
blackberry, evergreen	peavine, everlasting	birdsrape mustard	hogweed, giant
blackberry, Himalayan	ribbon grass	brome, ripgut	iris, yellow flag
canarygrass, reed	sowthistle, perennial	broom, Scotch	knapweed, diffuse
hawkweed, orange	tansy, common	burdock, common	knapweed, meadow
Hawkweed, yellow	thistle, Canada	butterfly bush	knapweed, spotted
knotweed, Bohemian	whitetop, hairy	carrot, wild	loosestrife, purple
knotweed, giant	yellow archangel	cheatgrass or downy brome	nightshade, hairy
knotweed, Japanese	-	chervil, wild	old man's beard
*Vegetative Propagation is the ability to reproduce from root and stem fragments. Indicated species must be thoroughly excavated from the ground, collected, and disposed into trash for control.		chicory	poison hemlock
		cinquefoil, sulfur	tansy, ragwort
		comfrey	teasel, common
		English hawthorn	thistle, bull
		fennel, common	wormwood, absinth

Control:

Do NOT mow any plants listed for control in table above

Completely remove all above ground biomass and as much of the root system as feasible (care must be taken to ensure proper collection and disposal of waste)

Ensure all flowers and seed heads are cut, bagged and disposed of into trash

Be aware of species listed as *Vegetative Propagation* and dispose of all plant material into trash and NOT compost

Additional control and identification resources can be found online at Clallam County's Noxious Weed page (<https://www.clallamcountywa.gov/821/Noxious-Weed-Control>) and the State Weed Board's website (<http://www.nwcb.wa.gov/>).

Sample Failure to Control Warning



Clallam County
223 E Fourth St, Suite 15
Port Angeles, WA 98362

Date

RE: Failure to fulfill 'Owner will Control' agreement

Dear Property Owner,

You entered into an Owner will Control agreement with Clallam County Road Department regarding noxious and invasive weeds on the county roadside adjacent to your property.

Crews were recently in your area and found the roadside adjacent to your property has not been maintained as required by the terms and conditions of the agreement (see enclosed).

You have ten (10) days from date of this letter to control weeds as outlined in the Owner Will Control agreement. If the right-of-way is not adequately maintained as described in the agreement the agreement will be immediately terminated and weeds of concern will be controlled as determined by the County, including by the use of herbicides.

If the Owner will Control agreement is terminated as described above, you may still apply to reenter into an Owner will Control agreement for next calendar year.

If you have any questions, please call _____ at _____

Or email us at: _____

This is the only notice you will receive regarding this matter.

Sincerely,

County Representative
Clallam County Roads Department

Enclosed: Owner Will Control agreement

APPENDIX G: ADOPT-A-PATCH PERMIT

Permits online at: <https://www.clallamcountywa.gov/821/Noxious-Weed-Control>

Clallam County Public Works Department
223 East Fourth Street, Suite 15 Port Angeles, WA 98362
360-417-2703 Phone 360-417-2414 Fax

\$160 plus all costs beyond public use**

**See C.C.C. 5.100.245 – Fee Schedule 245-A

PROJECT NO. _____
ROAD NAME _____
PERMIT NO. _____
COUNTY USE ONLY

APPLICATION FOR SPECIAL USE OR EVENT ALONG CLALLAM COUNTY RIGHT OF WAY

In Clallam County, a "Right-of-Way" permit is required to work along a county-owned road within the county right of way.

PLEASE PRINT

Name of Applicant: _____	County Road: _____
Mailing Address: _____ _____ _____	Address/ Milepost of Project Site: _____ When the project is approved: (check one item below) ☐ Mail permit when approved ☐ Call when approved ☐ Fax when approved
Phone: _____	
Cell Phone: _____	
Fax: _____	

USE PROPOSED & PURPOSE

Special Use: NOXIOUS WEED CONTROL

Name of Event Coordinator: _____

Start Date _____

End Date _____

IMPORTANT:

Project Location(s) Description: _____

(Reference "IWM Partner List" for available locations)

THE EXACT LOCATION OF THE ENTIRE EVENT/USE AREA MUST BE CLEARLY MARKED SO AS TO BE EVIDENT TO COUNTY PERSONNEL. FAILURE TO COMPLY WILL RESULT IN A DELAY OF THE PROCESSING OF THIS PERMIT.

It is the responsibility of the applicant to notify all utilities and private property owners when such property is liable to injury or damage through the performance of the permitted work. The applicant shall make all necessary arrangements relative to the protection of such property and/or utilities.

By signing this permit, the applicant agrees to comply with all conditions as stated on the PERMIT, Form RWPCOND041604, Permit Conditions Addendum and C.C.C. 5.100.245 – Fee Schedule 245-A. Applicant has 10 days from permit approval date to request clarification or modification to permit conditions attached.

Signed _____ Date _____

***** COUNTY USE ONLY *****

PERMISSION IS HEREBY ☐ GRANTED ☐ DENIED

Call 360-417-2703 for the following:

☒ Start Date ☐ _____ ☐ _____ ☒ Final

The Approved Permit Must be Posted on Site Until Final Inspection.

COMMENTS: _____

FEE CALCULATION

AMT WAIVED: _____

NET FEE: _____

DATE: _____

RECEIPT# _____

CHECK# _____

REC'D BY: _____

This permit shall be void unless the work herein contemplated is completed before the following date: _____

Area Supervisor/Design Review Engineer _____

Date _____

Final Inspection By: _____

Date: _____

APPENDIX H: ROADSIDE TREATMENT ACTIVITIES 2024 (TABLE 14)

This table includes all county roadsides managed for noxious weeds in 2024 under the Clallam County Road Department IWM Plan. The table is sorted alphabetically by road name. Definitions for the headings can be found at the end of the table. Species treated are listed alphabetically by the assigned 4-letter code (see Appendix B); 4-letter codes shown in bold are regulated noxious weeds and required for control in Clallam County.

We completed **310 treatments** on **198 county roads** over **95 days** and controlled **36 species**. In total, we treated **298.4 miles** (including retreatments/spot treatments) and examined **1,113.0 acres** (including monitoring surveys) of county roadside. For retreatments, Miles Examined, Acres Examined, and Acres Treated were counted in full in order to correctly calculate application rates and Solid Acres.

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
3 Crabs Rd	Herbicide	04/01/24	0.10	2.53	0.100	0.0115	0.000	COMA	2018-2022, 2024
Agate Beach Rd	Manual	08/20/24	0.20	0.50	0.100	0.0000	0.000	JAVU	2024
Aster Rd	Manual	07/18/24	0.10	0.25	0.250	0.0000	0.001	JAVU	2018-2021, 2023, 2024
	Manual	10/22/24	0.10	0.24	0.010	0.0000	0.000	JAVU	
Atterberry Rd	Manual	02/05/24	0.01	5.10	0.010	0.0000	0.000	COMA	2019, 2022, 2023, 2024
	Manual	09/03/24	0.10	6.78	0.240	0.0000	0.000	JAVU	
Baker Farm Rd	Manual	08/05/24	0.10	0.96	0.240	0.0000	0.000	ARMI	2024
Ballard Rd	Herbicide	04/22/24	0.20	0.35	0.350	0.0161	0.000	LAGA, CYSC, RUAR	2024
Beach Dr	Manual	05/28/24	0.50	1.00	1.000	0.0000	0.000	COMA	2024
	Manual	10/22/24	0.10	1.21	0.010	0.0000	0.000	COMA	
Bishop Rd	Manual	08/21/24	0.20	0.96	0.480	0.0000	0.000	CEMO	2024
Black Diamond Rd	Manual	06/18/24	4.40	10.60	1.270	0.0000	0.001	JAVU, DIFU	2017-2024
	Manual	08/21/24	0.10	10.66	0.240	0.0000	0.001	TAVU	
Blue Mountain Rd	Manual/Herbicide	08/26/24	7.00	16.96	16.960	0.0000	0.000	JAVU, POBO, HYPE, CIAR, CEMO, LALA	2024
	Manual	09/04/24	0.10	16.96	0.240	0.0459	0.000	CEMO	
Bogachiel Way	Manual	07/31/24	0.30	2.42	0.720	0.0000	0.004	JAVU	2024
Calawah Way	Manual	07/31/24	1.400	3.390	3.390	0.0000	0.004	JAVU	2023-2024
	Manual	08/12/24	1.400	3.390	3.390	0.0000	0.009	JAVU	

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Cameron Rd (District 2)	Manual	12/17/24	0.10	1.93	0.240	0.0000	0.0012	JAVU	2024
Camp Hayden Rd	Manual	08/20/24	3.50	8.48	8.480	0.0574	0.000	JAVU	2018-2024
	Herbicide	08/21/24	3.50	8.48	8.480	0.0000	0.001	GERO, CLVU, JAVU, CIVU	
	Manual	11/01/24	0.10	8.48	0.240	0.0000		JAVU	
Carlsborg Rd	Herbicide	03/29/24	1.80	3.26	3.260	0.0505	0.000	CEST, CYSC, CIAR, CIVU, TAVU, RUAR, CEMO	2018-2024
	Manual	09/04/24	0.10	4.36	0.240	0.0000	0.000	CEST	
Cassidy Rd	Herbicide	09/03/24	0.80	1.93	1.930	0.0138	0.000	JAVU, CEMO, CIAR, CIVU, DACA, DALA	2024
Cedar Park Dr	Manual	10/28/24	0.10	1.69	0.240	0.0000	0.0001	JAVU	2024
Chicken Coop Rd	Manual/Herbicide	07/06/24	3.50	8.48	8.480	0.0161	0.001	CIAR, CIVU, CLVU, DIFU, JAVU	2017-2020, 2024
Cline Spit Rd	Herbicide	04/17/24	0.20	0.36	0.360	0.0000	0.000	COMA	2021, 2023-2024
Coho Dr	Manual	09/11/24	0.10	0.72	0.240	0.0000	0.000	JAVU	2021, 2023-2024
Commercial St	Herbicide	07/10/24	0.10	0.24	0.240	0.0023	0.000	JAVU	2023-2024
	Manual	09/16/24	0.10	0.24	0.240	0.0000	0.000	JAVU	
Cooper Ranch Rd	Manual/Herbicide	08/27/24	4.20	10.20	10.200	0.0046	0.000	JAVU, CLVU, GERO, RULA	2019-2024
Coulter Rd	Herbicide	03/05/24	0.20	0.36	0.360	0.0057	0.000	COMA, CIVU, CIAR	2020-2022, 2024
	Herbicide	10/09/24	0.10	0.96	0.100	0.0000	0.000	COMA	
Crescent Beach Rd	Manual	08/20/24	1.50	8.48	3.630	0.0000	0.001	JAVU, CEMO	2020-2024
Dan Kelly Rd	Herbicide	07/17/24	3.20	7.80	7.800	0.0459	0.000	POBO, JAVU, CIVU, CIAR, DACA, CEMO, CEDI	2017-2021, 2023-2024
	Manual	08/20/24	3.20	7.75	7.750	0.0000	0.001	JAVU	
Deer Park Rd	Herbicide	03/29/24	0.10	0.10	0.100	0.0115	0.000	FOVU	2018-2024
	Herbicide	09/04/24	0.10	11.63	0.240	0.0023	0.000	CEMO	
	Manual	10/28/24	0.10	10.90	0.240	0.0000	0.0004	JAVU	

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Diamond Point Rd	Manual	10/22/24	1.00	9.68	2.420	0.0000	0.000	JAVU	2018-2024
Dietz Rd	Manual	08/05/24	0.50	1.21	1.210	0.0000	0.000	JAVU, CIVU	2021-2022, 2024
Discovery View Dr	Manual	10/22/24	0.10	1.21	0.010	0.0000	0.000	JAVU	2017, 2019, 2020, 2024
Division St	Herbicide	07/10/24	0.10	0.24	0.240	0.0046	0.000	JAVU	2024
Don Schmith Rd	Manual/Herbicide	09/18/24	0.50	1.21	1.210	0.0367	0.000	JAVU, CEMO, LALA, PORE, CIAR, RUAR	2023-2024
Doss Rd	Manual	12/18/24	0.10	1.21	0.240	0.0000	0.0001	JAVU	2024
Draper Rd	Manual	08/05/24	0.10	2.42	0.240	0.0000	0.000	JAVU	2024
Dry Creek Rd	Manual	09/17/24	0.10	1.96	0.240	0.0000	0.000	JAVU	2021, 2024
Durrwachter Rd	Manual	08/20/24	1.50	3.63	3.630	0.0000	0.002	JAVU	2019, 2024
Eagle Point Rd	Herbicide	09/16/24	0.20	3.15	0.480	0.0069	0.000	POBO	2024
East Beach Rd	Manual/Herbicide	04/30/24	1.40	2.53	2.530	0.0002	0.035	CYSC, DALA, GERO, DIPU, ANCA	2017-2024
East Lake Pleasant Rd	Manual	09/11/24	1.20	2.90	2.900	0.0000	0.001	JAVU	2024
East Lyre River Rd	Manual	10/10/24	0.60	1.45	1.450	0.0000	0.000	JAVU, CEMO	2017-2024
East Sequim Bay Rd	Manual/Herbicide	08/07/24	4.10	9.93	9.930	0.0000	0.000	CLVU, JAVU	2019-2022, 2024
	Manual	08/19/24	0.10	9.93	0.240	0.0046	0.000	JAVU	
	Manual/Herbicide	10/01/24	4.10	10.00	10.000	0.0115	0.000	CLVU, JAVU	
Easterly Rd	Herbicide	04/29/24	0.30	0.54	0.540	0.0230	0.000	CEMO, CYSC, RUAR, HYPE, GERO	2017-2021, 2023-2024
Eden Valley Rd	Manual	08/20/24	3.50	8.480	8.480	0.0000	0.002	JAVU, DIFU, CEMO	2017-2024
Elk Creek Ridge Rd	Manual/Herbicide	07/31/24	0.20	0.480	0.480	0.0023	0.001	JAVU	2024
Elk Loop Dr	Manual	07/31/24	0.20	0.480	0.480	0.0000	0.001	JAVU	2023-2024

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Elk Valley Rd	Manual/ Herbicide	08/12/24	0.50	1.210	1.210	0.0551	0.010	JAVU	2023- 2024
	Manual	10/30/24	0.10	1.210	0.240	0.0000	0.0001	JAVU	
Elwha River Rd	Manual	09/17/24	0.50	4.60	1.210	0.0000	0.000	JAVU	2019- 2024
Emery Rd	Manual	08/26/24	0.50	1.21	0.100	0.0000	0.000	CIVU	2024
Evans Rd	Manual	07/15/24	1.70	4.20	4.200	0.0000	0.002	DIFU	2018- 2021, 2024
Evergreen View Pkwy	Manual	08/20/24	0.50	1.21	1.210	0.0000	0.000	JAVU	2019, 2024
Farrington Rd	Herbicide	04/10/24	0.90	1.63	1.630	0.0069	0.000	JAVU, RUAR, CIAR, CIVU, CEMO	2017- 2024
	Manual	09/17/24	0.90	2.18	2.180	0.0000	0.002	JAVU, CEMO	
Fasola Rd	Manual	02/12/24	0.30	1.81	0.540	0.0023	0.000	DIFU	2019- 2022, 2024
	Herbicide	03/06/24	0.30	0.54	0.540	0.0207	0.000	DIFU	
	Herbicide	04/08/24	0.70	1.27	1.270	0.0000	0.001	DIFU, CIVU, RUAR, FOVU	
	Manual	07/16/24	1.00	2.42	2.420	0.0000	0.001	COMA, DIFU	
Fey Rd	Manual	04/10/24	0.30	0.540	0.540	0.0000	0.000	JAVU	2024
Fisher Cove Rd	Herbicide	10/10/24	0.80	1.93	1.930	0.0390	0.000	CEMO, CIVU, DIPU	2017- 2021, 2024
Fleming Dr	Manual	07/15/24	0.40	1.00	1.000	0.0000	0.000	JAVU	2017, 2019- 2020, 2024
	Manual	10/22/24	0.10	0.96	0.010	0.0000	0.000	JAVU	
Freshwater Bay Rd	Manual	08/20/24	1.00	6.06	2.420	0.0000	0.002	JAVU	2019- 2021, 2023- 2024
	Manual	11/01/24	2.50	6.06	6.060	0.0000	0.0002	JAVU	
Freshwater Park Lp	Manual	04/10/24	1.30	2.35	2.350	0.0000	0.000	JAVU	2022- 2024
	Manual	08/20/24	1.30	3.15	3.150	0.0000	0.001	JAVU	
	Manual	11/01/24	1.30	3.15	3.150	0.0000	0.0004	JAVU	
Front St	Manual/ Herbicide	07/10/24	0.60	1.45	1.450	0.0230	0.002	JAVU	2023- 2024
Gagnon Rd	Manual	07/22/24	0.10	1.210	0.240	0.0023	0.000	JAVU	2021, 2024
	Herbicide	09/17/24	0.10	1.210	0.240	0.0000	0.000	CEMO	
Gasman Rd	Herbicide	05/30/24	1.30	3.900	3.200	0.2984	0.000	CYSC, HYPE, CIVU, RUAR, DIPU, GERO	2017, 2020, 2024

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Gaydeski Rd	Manual	08/13/24	0.40	0.960	0.960	0.0000	0.000	JAVU	2024
Gilbert Rd	Herbicide	02/07/24	0.20	0.400	0.40	0.0115	0.000	COMA, DALA, CIVU, ANCA, CYSC	2020-2024
	Herbicide	06/05/24	0.10	0.400	0.10	0.0023	0.000	COMA, RUAR	
Glass Rd	Manual	08/05/24	2.20	5.330	5.33	0.0000	0.000	JAVU, CIVU	2021-2022, 2024
Gossett Rd	Manual	09/17/24	0.90	2.180	2.18	0.0000	0.000	JAVU, CEMO	2017-2024
	Manual/Herbicide	08/21/24	0.90	2.180	2.18	0.0092	0.000	CEMO, JAVU, GERO, CIAR, CIVU, RUAR	
Granite Rd	Herbicide	05/16/24	0.30	3.34	3.34	0.1768	0.000	CYSC, CEMO, HYPE, CIVU, CIAR, RUAR	2021-2022, 2024
Grant Rd-District 1	Manual	08/06/24	0.20	0.72	0.48	0.0000	0.000	CEST	2018, 2022, 2024
Grael-Ramapo Rd	Manual	08/21/24	1.90	4.60	4.60	0.0000	0.001	JAVU	2018-2021, 2023-2024
Happy Valley Rd	Manual	02/12/24	1.00	1.81	1.81	0.0849	0.000	DIFU	2017-2024
	Manual	02/27/24	3.30	6.00	0.20	0.0000	0.005	DIFU	
	Manual	03/27/24	0.10	0.10	0.10	0.0000	0.001	DIFU	
	Herbicide	04/29/24	2.00	14.44	14.44	0.0000	0.004	CEMO, CEST, JAVU, RUAR, CYSC, CIVU, CIAR	
	Manual/Herbicide	09/23/24	1.25	3.00	3.00	0.0861	0.002	CEST, CEMO, CIAR, DIFU	
	Manual	12/18/24	0.40	7.27	0.96	0.0000	0.0063	DIFU	
	Manual	12/20/24	0.20	6.00	0.50	0.0000	0.0035	CEMO	
Hardwick Rd	Manual	08/07/24	0.10	0.96	0.10	0.0000	0.000	JAVU	2024
Harrison Rd	Manual	10/22/24	0.30	0.72	0.72	0.0000	0.003	BEIN, CEST	2021-2022, 2024
Harry Brown Rd	Manual	06/18/24	0.20	0.40	0.40	0.0000	0.000	JAVU	2024
Hauk Rd	Herbicide	09/16/24	0.10	0.96	0.24	0.0161	0.000	POBO	2024
Henry Boyd Rd	Manual	08/05/24	0.50	3.87	1.21	0.0000	0.002	JAVU	2021-2024

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Herrick Rd	Herbicide	07/22/24	0.10	3.630	0.10	0.0023	0.000	CEST	2021-2022, 2024
Hoare Rd	Manual	06/18/24	0.10	1.30	0.10	0.0000	0.000	CYSC	2023-2024
Hoko-Ozette Rd	Manual/Herbicide	06/06/24	21.00	50.10 0	50.10	0.0046	0.000	POBO, JAVU, CYSC	2017-2024
	Herbicide	07/10/24	17.00	44.40 0	31.50	0.0000	0.006	CIVU, HIAU, POBO, RUAR	
	Manual	07/30/24	18.00	53.30 0	43.60	0.0115	0.000	JAVU	
Holgerson Rd	Manual	03/12/24	0.10	0.72	0.10	0.0000	0.001	DIFU	2024
	Manual	09/10/24	0.10	0.96	0.24	0.0000	0.001	DIFU	
Holland Rd	Manual	08/06/24	0.10	4.12	0.24	0.0000	0.000	JAVU	2018-2021, 2024
Hooker Rd	Herbicide	02/07/24	0.40	0.76	0.76	0.0230	0.000	COMA, LEVU, RUAR	2020-2022, 2024
	Herbicide	06/05/24	5.00	1.00	1.00	0.0689	0.000	HYPE, DIFU, RUAR, CIAR, CIVU	
Huckleberry Hill Dr	Manual	08/20/24	0.40	0.96	0.96	0.0000	0.001	JAVU	2019, 2024
	Manual	11/01/24	0.10	0.96	0.24	0.0000	0.000	JAVU	
Hudson Rd	Herbicide	04/10/24	0.30	0.54	0.54	0.0023	0.000	CIVU, CIAR, DIPU	2024
Industrial Pkwy	Manual	07/18/24	0.50	1.20	1.20	0.0000	0.000	JAVU	2017, 2020, 2024
James Page Rd	Herbicide	08/06/24	0.10	0.720	0.10	0.0184	0.000	CIAR	2024
Jamestown Rd	Manual	09/10/24	0.10	3.87	0.24	0.0000	0.000	FOVU	2019, 2024
Johnson Creek Rd	Herbicide	04/29/24	0.50	0.90	0.90	0.0230	0.000	CYSC, CEMO, RUAR, HYPE, JAVU, CIVU, CIAR	2017-2024
	Herbicide	09/23/24	0.25	1.50	0.60	0.0057	0.000	CEMO, CIAR	
Joyce-Piedmont Rd	Manual	08/21/24	0.10	3.39	0.10	0.0000	0.004	TAVU	2018-2021, 2024
Kemp St	Herbicide	09/16/24	0.10	0.100	0.10	0.0069	0.000	POBO	2024
Kirner Rd	Manual	08/06/24	0.10	1.210	0.24	0.0000	0.000	CEST	2024
	Manual	08/14/24	0.10	2.180	0.24	0.0000	0.000	TAVU	
Kitchen-Dick Rd	Herbicide	03/04/24	0.20	5.810	0.36	0.0115	0.000	COMA	2017-2024

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Lake Creek Rd	Manual	09/11/24	0.30	0.72	0.72	0.0000	0.000	JAVU	2024
Lamar Ln	Herbicide	03/20/24	0.40	1.270	0.72	0.0115	0.000	CAPY, COMA, CIVU, CIAR	2019-2021, 2023-2024
Lawrence Rd	Herbicide	04/10/24	0.70	1.270	1.27	0.0115	0.000	JAVU, CIVU, CIAR, CYSC	2024
Liljedahl Rd	Manual	08/21/24	1.00	2.420	2.42	0.0000	0.000	JAVU	2018-2019, 2023-2024
Little Loop Dr	Manual	08/05/24	0.20	1.930	0.48	0.0000	0.001	JAVU	2020-2024
Little River Rd	Herbicide	08/28/24	3.50	8.48	8.48	0.0597	0.000	CEMO, CLVU, CIVU, GERO, JAVU	2017-2024
Lost Mountain Rd	Manual	08/26/24	5.20	12.60	12.60	0.0000	0.000	JAVU	2019-2024
	Manual/Herbicide	09/30/24	3.40	8.24	8.24	0.0689	0.000	JAVU, CIAR, CIVU, DIFU, GERO, CIIN, CEMO	
Lotzgesell Rd	Herbicide	03/07/24	1.50	8.00	2.71	0.0402	0.000	COMA, DIFU, CYSC, CIVU	2018-2024
	Herbicide	03/12/24	0.01	0.01	0.01	0.0002	0.000	COMA	
	Manual	12/20/24	0.10	8.24	0.10	0.0000	0.0001	COMA	
Louella Rd	Herbicide	09/18/24	0.10	2.18	0.10	0.0023	0.000	CEMO	2024
Lower Elwha Rd	Manual	01/08/24	0.01	0.100	0.100	0.0000	0.008	CYSC	2019-2024
	Manual/Herbicide	07/22/24	2.80	6.780	6.780	0.0781	0.004	CIAR, CIVU, CEMO, CEST, DACA, JAVU	
	Manual	11/01/24	1.40	6.800	3.400	0.0000	0.0004	JAVU	
Lucken Dr	Manual	07/31/24	0.00	0.240	0.000	0.0000	0.000	JAVU, CIVU	2024
Lupine Dr	Manual	07/18/24	0.50	1.20	1.20	0.0000	0.000	JAVU	2017-2020, 2024
	Manual	10/22/24	0.10	1.21	0.24	0.0000	0.000	JAVU	
Madrona Way	Manual	07/15/24	1.10	2.700	2.70	0.0000	0.003	JAVU	2017, 2019-2020, 2022-2024
	Manual	10/22/24	0.10	2.700	0.24	0.0000	0.000	JAVU	
Manzanita Dr	Manual	05/28/24	0.7	1.30	1.30	0.0000	0.000	JAVU	2020, 2023-2024
	Manual	07/18/24	0.7	1.70	1.70	0.0000	0.000	JAVU	

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Mapleton Way	Manual	07/22/24	0.1	0.72	0.24	0.0000	0.000	JAVU	2023-2024
Marine Dr	Herbicide	03/20/24	0.10	1.450	0.10	0.0023	0.000	COMA	2019-2024
	Herbicide	04/17/24	0.10	3.800	0.10	0.0023	0.000	COMA	
Marmot Loop	Herbicide	04/08/24	0.10	0.20	0.10	0.0046	0.000	ANSY	2023-2024
Mary Clark Rd	Herbicide	08/27/24	7.70	18.60	18.60	0.0298	0.002	JAVU, RULA, HYPE	2019-2024
	Manual	09/11/24	7.70	18.66	18.66	0.0000	0.000	JAVU	
	Manual	12/16/24	0.10	18.60	0.24	0.0000	0.0001	JAVU	
Maxfield Rd	Manual	08/13/24	0.80	1.930	1.93	0.0000	0.000	JAVU	2024
McAlmond St	Herbicide	04/17/24	0.10	0.240	0.24	0.0344	0.000	CYSC	2024
McGarvie Rd	Manual	09/17/24	0.30	0.720	0.72	0.0000	0.000	JAVU	2017-2019, 2023-2024
Melton Rd	Manual	06/18/24	0.10	0.540	0.10	0.0000	0.000	HIAU	2023-2024
Merchants Rd	Manual	07/31/24	0.50	1.210	1.21	0.0000	0.000	JAVU	2024
Mina Smith Rd	Manual/Herbicide	04/22/24	3.30	8.00	8.00	0.0344	0.002	JAVU, LAGA, CYSC, CIVU	2018-2024
	Manual/Herbicide	07/02/24	1.55	8.00	4.10	0.0023	0.005	CYSC, LAGA	
	Manual	10/25/24	1.10	8.00	2.66	0.0000	0.0002	JAVU	
Monroe Rd	Manual	08/05/24	0.60	9.21	1.45	0.0000	0.001	JAVU, CEMO	2018, 2020, 2023-2024
Moriarty Rd	Manual	10/25/24	0.10	3.15	0.100	0.0000	0.0000	JAVU	2024
Mount Angeles Rd (South of Eckard Rd)	Manual	08/19/24	0.10	10.66	0.240	0.0046	0.000	JAVU	2024
	Herbicide	08/28/24	0.10	0.10	0.100	0.0023	0.000	POBO	
	Herbicide	09/30/24		0.10	0.100	0.0000	0.000	HIAU	
Mount Pleasant Rd	Manual	01/04/24	0.10	0.36	0.18	0.0689	0.000	CYSC	2020-2024
	Manual	01/05/24	1.50	2.00	1.00	0.0046	0.000	CYSC	
	Herbicide	03/12/24	1.60	5.43	2.90	0.0023	0.000	COMA, CEMO	
	Herbicide	07/11/24	0.10	0.20	0.20	0.0000	0.019	TAVU	
	Herbicide	07/17/24	0.10	0.10	0.10	0.0000	0.010	TAVU	

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
	Manual	08/05/24	1.50	13.80	3.63	0.0000	0.001	CIVU, JAVU	
Myrtle St	Herbicide	09/30/24	0.10	0.10	0.100	0.0002	0.000	TAVU	2024
Nichols Rd	Manual	06/06/24	0.60	1.10	1.100	0.0000	0.000	CIVU	2024
North Bay View St	Herbicide	04/12/24	0.10	0.60	0.600	0.0459	0.000	TAVU, CYSC, CIVU	2024
North Brown Rd	Manual	09/11/24	0.10	1.20	0.100	0.0000	0.001	DIFU	2020, 2022 2024
North Diamond Shore Ln	Manual	05/28/24	0.10	0.72	0.100	0.0000	0.000	GERO	2024
North Eldridge Rd	Manual	09/03/24	0.10	0.48	0.100	0.0000	0.000	JAVU	2024
North Masters Rd	Herbicide	04/19/24	0.13	1.10	1.100	0.0918	0.000	CYSC	2018, 2024
	Herbicide	04/24/24	0.13	1.10	1.100	0.0505	0.000	CYSC, RUAR, CIVU	
North St	Manual	05/28/24	0.40	0.72	0.010	0.0000	0.000	JAVU	2019-2020, 2024
	Manual	10/22/24	0.10	0.96	0.100	0.0000	0.002	JAVU	
O'Brien Rd	Manual	12/19/24	0.20	9.93	0.200	0.0000	0.0039	CEST, PORE	2017-2021, 2023-2024
Okerman Rd	Manual	10/15/24	0.40	0.96	0.960	0.0000	0.001	CEMO	2020, 2021, 2023-2024
Old Black Diamond	Manual	06/18/24	0.20	0.40	0.400	0.0000	0.000	JAVU	2021, 2023-2024
Old Blyn Hwy	Herbicide	02/08/24	0.50	2.36	0.900	0.0230	0.000	COMA, JAVU	2018-2024
	Herbicide	08/14/24	2.10	5.09	5.090	0.0482	0.000	CLVU, JAVU, CIAR, CIVU, GERO, COMA	
	Herbicide	10/02/24	0.30	0.72	0.720	0.0758	0.000	CIAR, CLVU, GERO, RUAR	
	Manual	10/28/24	0.10	5.09	0.240	0.0000	0.0004	COMA	
Old Mill Rd (South of 4704 address)	Herbicide	09/09/24	0.90	2.18	2.180	0.0046	0.000	CEMO, POBO	2024

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Old Olympic Hwy	Manual	03/27/24	0.10	0.10	0.100	0.0000	0.003	DIFU	2017-2024
	Manual/Herbicide	07/11/24	0.10	3.87	0.100	0.0000	0.001	CEST, COMA	
	Manual	07/18/24	1.20	2.90	2.900	0.0000	0.008	JAVU	
	Manual	07/18/24	0.20	2.00	2.000	0.0000	0.000	DIFU	
	Manual	08/06/24	0.10	4.84	0.240	0.0000	0.000	FOVU, CEST	
	Manual	10/22/24	0.10	0.10	0.100	0.0046	0.000	FOVU	
	Manual	10/28/24	0.20	0.48	0.480	0.0000	0.0002	DIFU	
Olympic Hot Springs Rd	Herbicide	08/28/24	2.00	4.84	4.840	0.0803	0.000	CEMO, GERO, CLVU, CIAR, CIVU, ARMI, JAVE, POBO, LALA	2017-2021, 2023-2024
Palmer Rd	Manual	07/31/24	0.10	1.21	0.240	0.0000	0.000	JAVU	2024
Palo Alto Rd	Herbicide	03/05/24	0.10	0.10	0.100	0.0002	0.000	COMA, DACA	2017-2024
	Manual/Herbicide	09/18/24	8.70	21.09	21.090	0.0551	0.000	JAVU, CEMO, CIAR, PORE, LEVU, CLVU	
Panorama Blvd	Manual	04/02/24	0.30	1.00	0.400	0.0000	0.001	JAVU	2019-2021, 2024
Pavel Rd	Herbicide	08/27/24	0.10	3.15	0.100	0.0023	0.000	POBO	2019, 2024
Pit Ln	Manual	06/25/24	0.90	2.20	2.200	0.0000	0.001	JAVU	2023-2024
Place Rd	Manual	08/20/24	2.50	6.06	6.060	0.0000	0.000	JAVU	2021-2022, 2024
Poplar Ct	Manual	05/28/24	0.10	0.18	0.100	0.0000	0.000	JAVU	2017, 2020, 2024
	Manual	10/22/24	0.10	0.24	0.240	0.0000	0.000	JAVU	
Port Williams Rd	Manual	07/15/24	0.10	5.57	0.100	0.0000	0.000	COMA	2018, 2020-2022, 2024
Prawn Rd	Manual	08/20/24	0.10	0.48	0.240	0.0000	0.000	CIVU	2024
Priest Rd	Manual	08/06/24	0.10	0.72	0.240	0.0000	0.000	CEMO	2024
Quillayute Rd	Manual/Herbicide	06/17/24	6.90	16.70	16.700	0.0000	0.004	CYSC, JAVU	2020-2024
	Manual	09/11/24	2.00	4.84	4.840	0.0069	0.005	JAVU	
	Manual	10/25/24	3.00	16.72	7.270	0.0000	0.0007	JAVU	

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Quillayute Airport Rd	Manual	10/25/24	0.40	0.96	0.960	0.0000	0.0002	JAVU	2021-2024
Reservoir Rd	Manual	06/18/24	0.20	0.40	0.100	0.0000	0.000	JAVU	2024
Rhododendron Dr	Manual	05/28/24	0.80	1.50	1.500	0.0000	0.001	JAVU	2017, 2019-2020, 2022, 2024
	Manual	07/18/24	0.90	2.20	2.200	0.0000	0.001	JAVU	
	Manual	10/22/24	0.10	1.93	0.010	0.0000	0.000	JAVU	
Rice St	Herbicide	07/10/24	0.10	0.24	0.240	0.0069	0.000	HIAU, JAVU	2024
Richwine Rd	Manual	09/11/24	0.60	1.45	1.450	0.0000	0.000	JAVU	2021-2024
Ridge View Dr	Herbicide	03/29/24	0.10	1.45	0.100	0.0115	0.000	COMA	2023-2024
River Rd	Herbicide	09/23/24	1.00	2.40	2.400	0.0689	0.000	CEMO, CIAR, CIVU, RUAR, GERO, PORE	2017-2021, 2023-2024
Scenic Overlook Way	Manual	10/28/24	0.10	0.24	0.240	0.0000	0.0000	FOVU	2024
Schindler Rd	Manual/Herbicide	02/08/24	0.01	0.54	0.010	0.0002	0.000	COMA, JAVU	2024
Schmitt Rd	Manual	09/17/24	0.50	1.21	1.210	0.0000	0.001	JAVU	2017-2021, 2023-2024
Schmuck Rd	Manual	07/15/24	0.20	3.15	1.000	0.0000	0.001	DIFU	2018, 2020, 2024
Schoolhouse Point Ln	Manual	08/06/24	0.40	0.96	0.960	0.0000	0.000	JAVU	2024
Secor Rd	Manual	08/26/24	0.10	1.21	0.100	0.0000	0.000	CEST	2024
Sekiu River Rd	Manual/Herbicide	07/10/24	0.70	1.69	1.690	0.0115	0.000	CIVU, LAGA, POBO	2019, 2023-2024
	Manual/Herbicide	09/16/24	0.70	1.69	1.690	0.0069	0.000	POBO, RUAR, CIVU	
Senz Rd	Herbicide	09/30/24	0.10	1.21	0.240	0.0046	0.000	LALA	2024
Sequim-Dungeness Way (North of 1323 address)	Herbicide	02/07/24	0.10	0.20	0.200	0.0057	0.000	COMA	2017-2021, 2023-2024
	Herbicide	03/06/24	0.10	0.36	0.180	0.0046	0.000	COMA, CIVU	
	Herbicide	04/19/24	1.20	2.90	2.900	0.0344	0.000	CEST, CIVU, RUAR, CIAR, TAVU	
	Manual	05/28/24	0.10	5.40	0.100	0.0275	0.000	COMA	

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
	Manual	08/14/24	0.10	9.93	0.240	0.0000	0.000	JAVU	
	Manual	08/22/24	1.00	2.42	2.420	0.0000	0.001	DIFU	
	Herbicide	10/01/24	0.10	0.20	0.200	0.0000	0.003	CIAR, COAR, GERO, RUAR	
	Manual	10/28/24	0.10	0.24	0.240	0.0000	0.0001	DIFU	
Shadow Ln	Manual	08/21/24	0.40	0.96	0.960	0.0000	0.000	JAVU	2024
Sherwood Rd	Manual	07/15/24	0.30	0.70	0.700	0.0000	0.001	JAVU	2017-2020, 2022, 2024
Siebert Creek Rd	Manual	12/19/24	0.10	1.93	0.200	0.0000	0.0014	PORE	2019, 2024
South 3rd Ave (Reservoir Rd to Happy Valley Rd)	Herbicide	04/01/24	0.50	0.90	0.900	0.0184	0.000	CEMO, EULA, RUAR, JAVU, CIAR, CIVU, HYPE	2021, 2024
South Bagley Creek Rd	Herbicide	03/04/24	0.30	2.72	0.540	0.0115	0.000	COMA, CIVU, GERO, CYSC	2018-2020, 2024
South Shore Rd	Herbicide	08/21/24	0.50	5.09	1.210	0.0459	0.000	GERO, CLVU, HIAU	2018-2019, 2024
Sporseen Rd	Manual	12/18/24	0.10	2.18	0.240	0.0000	0.0001	DIFU	2024
Steelhead Ave	Manual/ Herbicide	08/13/24	0.10	1.69	0.100	0.0023	0.000	HIAU	2024
Swan Bay Rd	Herbicide	06/06/24	0.80	1.50	1.500	0.0023	0.000	JAVU	2018-2021, 2024
	Manual	07/30/24	0.40	1.93	0.960	0.0000	0.000	JAVU	
Taylor Ranch Rd	Herbicide	02/07/24	0.80	1.45	1.450	0.0230	0.000	COMA, GERO, JAVU, CYSC, LEVU, RUAR	2019-2020, 2024
	Herbicide	05/23/24	0.40	1.50	0.720	0.0115	0.000	COMA, CIVU	
	Manual	08/06/24	0.30	1.93	0.720	0.0000	0.001	CEMO, COMA	
	Manual	12/17/24	0.10	1.93	0.240	0.0000	0.0000	COMA	
Taylor St	Manual	09/11/24	0.10	0.24	0.240	0.0000	0.000	JAVU, HIAU	2024
Thompson Rd	Manual	08/19/24	1.10	2.66	2.660	0.0000	0.001	JAVU, TAVU	2024
Timber Rd	Manual	05/28/24	0.10	0.18	0.100	0.0000	0.000	JAVU	2019-2020, 2024
	Manual	07/18/24	0.10	0.25	0.250	0.0000	0.000	JAVU	

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Towne Rd	Herbicide	03/13/24	0.50	2.53	0.720	0.0230	0.000	COMA, DIFU, CIVU, RUAR, SIAR	2018-2024
	Manual	05/28/24	0.10	0.10	0.100	0.0000	0.001	DIFU	
	Manual	06/27/24	0.00	0.10	0.100	0.0000	0.001	DIFU	
	Manual	10/22/24	0.10	0.24	0.240	0.0000	0.001	FOVU	
	Herbicide	12/11/24	0.75	1.80	1.800	0.0344	0.0000	BRSP, CIAR, CIVU	
Trout Ave	Manual	08/13/24	0.50	1.21	1.210	0.0000	0.001	JAVU	2023-2024
Turnstone Ln	Herbicide	03/29/24	0.70	1.27	1.270	0.0184	0.000	CEST, DIPU, CYSC, RUAR, DALA, CIAR	2017-2024
	Herbicide	10/15/24	0.30	1.69	0.720	0.0275	0.000	CEST	
Twin View Dr	Herbicide	04/17/24	0.60	1.09	1.090	0.0023	0.000	CEMO	2024
Undi Rd	Herbicide	03/18/24	0.40	3.62	0.720	0.2984	0.000	RUAR, JAVU, CIVU, LEVU, CYSC	2024
	Herbicide	10/07/24	0.70	4.80	1.700	0.5969	0.000	RUAR, JAVU, CIVU, CIAR, CYSC, RULA	
Valley Center Pl	Herbicide	06/05/24	0.10	3.50	0.100	0.0002	0.000	CEST	2020-2021, 2024
Ward Rd	Herbicide	03/06/24	1.60	3.08	2.900	0.0023	0.000	COMA, DIFU	2018-2022, 2024
	Herbicide	04/15/24	0.40	0.72	0.720	0.0069	0.000	COMA, RUAR, CIVU	
	Manual	07/16/24	0.10	4.12	0.100	0.0000	0.000	COMA	
Wasankari Rd	Manual	08/21/24	1.00	2.42	2.420	0.0000	0.000	JAVU	2018, 2024
Washington Harbor Rd	Herbicide	03/05/24	0.80	2.53	1.450	0.0161	0.000	COMA, CIAR, CIVU, CYSC	2019, 2022, 2024
	Herbicide	10/09/24	0.00	0.20	0.200	0.0069	0.000	COMA, CIAR, CIVU	
Washington St	Herbicide	07/10/24	0.40	0.96	0.960	0.0161	0.000	POBO, GERO	2022, 2024
	Manual	09/16/24	0.10	0.96	0.240	0.0000	0.002	JAVU	
West Arnette Rd	Manual	02/27/24	0.01	0.18	0.100	0.0000	0.000	DIFU	2024
	Manual	08/05/24	0.10	0.20	0.200	0.0000	0.000	DIFU	
West Bluff Dr	Manual	09/11/24	0.10	0.72	0.010	0.0000	0.000	JAVU	2024
West Duvall Rd	Manual	09/18/24	0.20	0.48	0.480	0.0000	0.000	JAVU	2023-2024
West Edgewood Dr	Manual/Herbicide	09/17/24	1.50	4.84	3.630	0.0161	0.001	CEMO, JAVU, RUAR, LAGA, CIAR, CIVU	2017-2022, 2024

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
West Hendrickson Rd	Herbicide	03/13/24	0.60	1.09	1.090	0.0115	0.000	COMA, DIFU, SIAR, CIVU, GERO	2018, 2020, 2022, 2024
West Lyre River Rd	Herbicide	10/10/24	0.60	1.45	1.450	0.0115	0.000	CEMO, JAVU	2017-2024
West Runnion Rd	Manual/Herbicide	08/06/24	0.20	1.21	0.480	0.0046	0.000	CEST	2024
West Silberhorn Rd	Manual	10/22/24	0.10	0.50	0.010	0.0000	0.000	FOVU	2024
West Washington St	Herbicide	02/07/24	0.30	3.30	3.300	0.0057	0.000	DALA, CEST	2017-2024
	Herbicide	04/01/24	0.42	5.70	5.700	0.1722	0.000	CEST, HYPE, DIPU, RUAR	
	Herbicide	06/24/24	0.30	5.70	5.700	0.0689	0.000	CEST, HYPE, SOAU	
	Manual	08/06/24	0.20	1.00	1.000	0.0631	0.000	CEST	
	Herbicide	10/15/24	0.40	4.96	4.960	0.0000	0.001	CEST, RUAR, HYPE, PORE	
Whiskey Creek Beach Rd (Hwy 112 to Schmitt Rd)	Manual	09/17/24	0.50	1.21	1.210	0.0000	0.000	JAVU	2017-2024
Whitcomb-Diimmel Rd	Manual	06/25/24	1.50	3.70	3.700	0.0000	0.004	JAVU	2020-2024
	Manual	08/13/24	1.50	3.63	3.630	0.0000	0.006	JAVU, CIVU	
Wilson Rd	Herbicide	03/19/24	0.80	1.45	1.450	0.0230	0.000	CIVU, CYSC, JAVU, DIPU, HEHE, GERO	2018-2019, 2021, 2024
	Herbicide	08/27/24	0.80	1.93	1.930	0.0046	0.000	JAVU, CIVU	
Woodcock Rd	Manual	01/24/24	0.01	0.01	0.010	0.0459	0.000	DIFU	2017-2024
	Herbicide	03/20/24	0.80	1.45	1.450	0.1148	0.000	COMA, CIAR, CIVU, LEVU	
	Herbicide	04/18/24	4.50	10.90	10.900	0.0000	0.000	COMA, CYSC, CIVU, CIAR, DIFU, LALA, FOVU, DACA, CEMO, CEST, ANSY, RUAR, EULA, GERO, LUAR	
	Manual	06/24/24	0.10	0.10	0.100	0.0000	0.000	COMA	

ROAD NAME	TREATMENT METHOD ¹	TREATMENT DATE	MILES TREATED	ACRES EXAMINED ²	ACRES TREATED ³	SOLID CHEMICAL ACRES TREATED ⁴	SOLID MANUAL ACRES TREATED ⁵	TREATED SPECIES LIST ⁶	YEARS TREATED
Woods Rd	Manual/Herbicide	08/19/24	3.20	10.18	8.180	0.4213	0.000	JAVU, CLVU, TAVU, CIAR, GERO, HYPE	2018-2024
	Manual/Herbicide	08/19/24	1.00	10.18	2.000	0.0574	0.000	JAVU, CLVU, CIAR, GERO, HYPE	
Wye Rd	Manual	08/21/24	0.60	1.45	1.450	0.0000	0.000	JAVU	2019-2024
Youngquist Rd	Manual	09/18/24	0.30	2.18	0.720	0.0000	0.000	DIFU	2024
Zaccardo Rd	Manual	08/07/24	0.10	1.00	0.100	0.0000	0.000	JAVU	2024
Total Roads: 198	310 Treatments	95 Days of Treating	298.4 Miles	1113.0 Acres Examined	242 Acres Treated	4.7121 Acres Treated	0.2993 Acres Treated	36 Species Treated	

*Non-priority species treated intermittently, meaning the entire population was not controlled during treatment

¹**M** – Manual control; **H** – Chemical control; **M/H** – Combination of manual and chemical control

²**Examined Acres** – The total area searched for noxious weeds while crew was involved in treatment activities

³**Treated Acres** – The gross area encompassing all treatments per road per day

⁴**Solid Chemical Treated Acres** – The estimated net area if the plants were “clumped” together; calculated using the tank mix volume applied and calibrated sprayer output

⁵**Solid Manual Acres** – The estimated net area controlled by any manual means (pulling, digging, cutting, etc.) if the plants were “clumped” together; calculated by number of plants removed

⁶**Species Treated** – The 4-Letter Weed codes correspond to the species’ scientific name and can be found in Appendix B. Bolded species are regulated noxious weeds required for control in Clallam County

APPENDIX I: REFERENCES

Noxious weed lists

Washington State Noxious Weed List. WAC 16-750

<https://app.leg.wa.gov/wac/default.aspx?cite=16-750>

Clallam County Noxious Weed List and Supplement.

<https://www.clallamcountywa.gov/887/Weed-Information-Weed-Lists>

Pictures of Class A Noxious Weeds. Washington State Noxious Weed Control Board

<https://www.nwcb.wa.gov/class-a-noxious-weeds>

Pictures of Class B Noxious Weeds. Washington State Noxious Weed Control Board

<https://www.nwcb.wa.gov/class-b-noxious-weeds>

Pictures of Class C Noxious Weeds. Washington State Noxious Weed Control Board.

<https://www.nwcb.wa.gov/class-c-noxious-weeds>

Pictures of noxious weeds of Washington – monitor list. Washington State Noxious Weed Control Board.

<https://www.nwcb.wa.gov/noxious-weed-monitor-list>

Pictures of noxious weeds of Washington – quarantine list. Washington State Noxious Weed Control Board.

<https://www.nwcb.wa.gov/noxious-weed-quarantine-list>

Clallam County Noxious Weed Control Board.

<https://www.clallamcountywa.gov/821/Noxious-Weed-Control>

Publications of the Washington State Noxious Weed Control Board.

<https://www.nwcb.wa.gov/publications>

Literature

Bramble, W.C., W.R. Byrnes, and R.J. Hutnik. 1987. Development of plant cover diversity on an electric transmission right-of-way. pg. 89-93. In: Byrnes, W.R. and H.A. Holt (eds). Proceeding, Fourth Symposium on Environmental Concerns in Right-of-Way Management. Indianapolis, IN. Dept. Forestry & Nat. Res., Purdue University, W. Lafayette, IN. 595 pp.

Bramble, W.C., R.H. Yahner, and W.R. Byrnes. 1999. Effect of herbicide maintenance of an electric transmission line right-of-way on butterfly populations. Jour. Arboriculture 25(6):302-310.

Cal-IPC. 2015. Best Management Practices for Wildland Stewardship: Protecting Wildlife When Using Herbicides for Invasive Plant Management. Cal-IPC Publication 2015-1. California Invasive Plant Council, Berkeley, CA. Available: www.cal-ipc.org

Colquhoun, J. 2003. Pacific Northwest's least wanted list: Invasive weed identification and management. EC 1563, Oregon State University, Corvallis. 44 pp.

Durkin, P. R. 2014. Scoping/Screening Level Risk Assessment on Fluazifop-P-butyl FINAL REPORT. Syracuse Environmental Research Associates, Inc., 293.

Durkin, P, Follansbee, M. 2004. Imazapic- Human Health and Ecological Risk Assessment- Final Report. Syracuse Environmental Research Associates, Inc, Fayetteville, New York 13066. 110 pp.

DiTomaso, J.M., G.B. Kyser et al. 2013. Weed control in natural areas in the western United States. Weed Research and Information Center, University of California, Davis. 544 pp.

Harper-Lore, B.L., M. Johnson, and W.F. Ostrum. 2013 Vegetation Management: An Ecoregional Approach. US DOT, Fed. Highway Admin. 239 pg.

- Harper-Lore, B.L., M. Johnson, and M.W. Skinner (editors). 2007. Roadside weed management. FHWA-HEP-07-017, Fed. Highway Admin., US Dept. Transportation. 369 pp.
- Hill, K. and R. Horner. 2005. Assessment of alternatives in roadside vegetation management. Final Research Report, Washington State Transportation Center, University of Washington, Seattle. Agreement T2695, Task 67. <http://www.wsdot.wa.gov/NR/rdonlyres/0CB59701-542E-4DF2-B8C8-1ACA3CB72172/0/FinalUWReport.pdf>
- Holt, H.A., R.L. McKenzie, F. Whitford, H.L. Hipkins, J. DiTomaso, R. Dickens, J. Orr, G. Blase, P. Northcutt, and J. McKenzie. 2002. Model Certification Manual for Right-of-Way Pesticide Applicators. <http://www.agriculture.purdue.edu/pesp/rowmanual/>
- Hopwood, J. et.al. 2015. Literature review: Pollinator habitat enhancement and best management practice in highway rights-of-way. Fed. Highway Admin. https://www.environment.fhwa.dot.gov/ecosystems/documents/pollinators_BMPs_in_highway_ROW.pdf
- Hutchings, J., J.P. Rutan, J. Gollen, K. Mankle, and L. Baldwin. 2015. Whatcom County Integrated Vegetation Management Plan. <http://www.co.whatcom.wa.us/DocumentCenter/Home/View/11074>
- Kitsap County Noxious Control Weed Board. 2015. City of Bremerton Integrated Vegetation Management Plan. 75 pg. <http://www.ci.bremerton.wa.us/DocumentCenter/View/2277>
- Miller, T. W., C. Lucero. 2014. Meadow knapweed (*Centaurea debeauxii*) Response to Herbicide and Mechanical Control. Invasive Plant Science and Management 2014 7:503-510.
- Monaco, T.J., S.C. Weller and F.M. Ashton. 2002. Weed Science Principles and Practices, 4th Edition. John Wiley & Sons, Inc., New York. 671 pg.
- National Archive and Records Administration: Federal Register Volume 78, No 192, Thursday October 3, 2013/ Rules and Regulations. Endangered and Threatened Wildlife and Plants ; Designation of Critical Habitat for Taylor's Checkerspot butterfly and Streaked Horned Lark.
- Panther, K.E., M.H. Ralphs et.al. 2011. Plants poisonous to livestock in the western states. USDA, Agri. Bull. No. 415. 107 pp.
- Pimentel, P, S. McNair et al. 2001. Economic and environmental threats of alien plant, animal, and microbe invasions. Agriculture, Ecosystems and Environment 84 (2001) 1–20
- Prather, T. 2015. Section V. Non-cropland and right-of-way. In: Peachey, Ed (editor). Pacific Northwest Weed Management Handbook. Oregon State University, Corvallis. p. V-1-V-10.
- Prather, T., T. Miller, and A. Hulting. 2015. Section W: Control of problem weeds. In: Peachey, Ed (editor). Pacific Northwest Weed Management Handbook. Oregon State University, Corvallis. p. W-1-W-44.
- Shaner, D.L., I. Burke et al. (editors). 2014. Herbicide Handbook, 10th Edition. Weed Science Society of America, Lawrence, KS. 513 pg.
- Tu, M, C. Hurd and J.M. Randall. 2001. Weed Control Methods Handbook. The Nature Conservancy. (<http://www.invasive.org/gist/products/handbook/methods-handbook.pdf>)
- USDA Forest Service. 2005. Pacific Northwest Region Invasive Plant Program, Final Impact Statement Vol. I 173 pp, Vol. III Appendix B-S.
- USDA Forest Service. 2015. Pacific Northwest Research Station, Herbicides an unexpected ally for native plants in the war against invasive species. Science Findings: Issue 176: 1-2.
- USDA Pesticide-Use Risk Assessments - <http://www.fs.fed.us/foresthealth/pesticide/risk.shtml>
- USDA Risk Assessment Worksheets - <http://www.fs.fed.us/foresthealth/pesticide/worksheets.shtml>
- USDI BLM. 2007. Final Vegetation Treatments Using Herbicides Programmatic Environmental Impact Statement. BLM PEIS 2007: http://www.blm.gov/wo/st/en/prog/more/veg_eis.html

USDI BLM. 2015. Draft programmatic environmental impact statement for vegetation treatments using aminopyralid, fluroxypyr, and rimsulfuron on Bureau of Land Management lands in 17 western states.http://www.blm.gov/style/medialib/blm/wo/Planning_and_Renewable_Resources/vegeis.Par.79259.File.dat/Report%20Cover%20Biological%20Assessment%20for%20Vegetation%20using%20Three%20Herbicides.pdf

USDT Federal Highway Administration. 2005. The Nature of Roadsides and the Tools to Work with It. FHWA Pub. No. FHWA-EP-03-005.

USDT Federal Highway Administration. 2016. Pollinators and Roadsides: Best Management Practices for Managers and Decision Makers. Available:

https://www.environment.fhwa.dot.gov/ecosystems/Pollinators_Roadsides/BMPs_pollinators_roadsides.asp

Washington Natural Heritage Program. Geographic information system. Rare plants and high quality ecosystems. Wnhp data set July 2015.

Washington State Department of Transportation. Undated. Aminopyralid, Roadside Vegetation Management Herbicide Fact Sheet. 4 pg. <http://www.wsdot.wa.gov/NR/rdonlyres/CD6D91AC-D382-4E5D-AF1E-50C52C34484F/0/Aminopyralid.pdf>

Washington State Department of Transportation. 2003. 2,4-D, Roadside Vegetation Management Herbicide Fact Sheet. 4 pg. http://www.wsdot.wa.gov/NR/rdonlyres/C6CBCC5B-64F8-4017-A209-9C6116EB36DB/0/2_4D.pdf

Washington State Department of Transportation. 2006. Clopyralid, Roadside Vegetation Management Herbicide Fact Sheet. 4 pg. <http://www.wsdot.wa.gov/NR/rdonlyres/C3ECE3F9-BFCF-432D-ACB2-6C716BB8FADC/0/Clopyralid.pdf>

Washington State Department of Transportation. 2006. Glyphosate, Roadside Vegetation Management Herbicide Fact Sheet. 4 pg. <http://www.wsdot.wa.gov/NR/rdonlyres/A72C98BF-88CD-4BAA-9B0F-5BB709A0C564/0/glyphosate.pdf>

Washington State Department of Transportation. 2006. Imazapyr, Roadside Vegetation Management Herbicide Fact Sheet. 4 pg. <http://www.wsdot.wa.gov/NR/rdonlyres/C8EB1611-2699-48FB-B5B8-161D1223BC92/0/imazapyr.pdf>

Washington State Department of Transportation. 2006. Triclopyr, Roadside Vegetation Management Herbicide Fact Sheet. 4 pg. <http://www.wsdot.wa.gov/NR/rdonlyres/DC50D1A8-3711-4B5F-BF12-05912616A486/0/triclopyr.pdf>

Washington State Department of Transportation. 2013. Chapter 6, Roadside Management. Washington State Department of Transportation Maintenance Manual M 51-01.05. 34 pg. <http://www.wsdot.wa.gov/publications/manuals/fulltext/M51-01/Chapter6.pdf>

Washington State Department of Transportation. 2014. Northwest Region, Area 1 Integrated Roadside Vegetation Management Plan. http://www.wsdot.wa.gov/NR/rdonlyres/586B5B84-EB29-45E1-AA6E-EC620BB3CD55/0/NWarea1_2014combined.pdf

Washington State Department of Transportation. 2014. Olympic Region, Area 3 Integrated Roadside Vegetation Management Plan. http://www.wsdot.wa.gov/NR/rdonlyres/315D3EA0-A71F-44D5-92F4-8F494182BD68/0/OLarea3_IVMplan2014combined.pdf

Washington State Department of Transportation. 2014. Southwest Region, Area 3 Integrated Roadside Vegetation Management Plan. http://www.wsdot.wa.gov/NR/rdonlyres/28AD03A3-7994-4A84-8602-1B14AC68264F/0/SWarea3_2014combined.pdf

Westbrook, R.G. 1998. Invasive plants, changing the landscape of America: Fact book. Federal Interagency Committee for the Management of Noxious and Exotic Weeds (FICMNEW), Washington, DC. 109 pp.

Whitson, T.D., L.C. Burrill et al. 2006. Weeds of the West. 9th ed. Western Society of Weed Science, Las Cruces, NM. 628 pp.

Xerces Society. Pollinator-Conservation-Roadsides <http://www.xerces.org/pollinator-conservation-roadsides/>

Yahner, R.H. and R.J. Hutnik. 2005. Plant species richness on an electric transmission right-of-way using integrated vegetation management. *Jour. Arboriculture* 31(3):124-130.